

EXHIBIT 18

Endocrine Treatment of Gender-Dysphoric/ Gender-Incongruent Persons: An Endocrine Society* Clinical Practice Guideline

Wylie C. Hembree,¹ Peggy T. Cohen-Kettenis,² Louis Gooren,³ Sabine E. Hannema,⁴ Walter J. Meyer,⁵ M. Hassan Murad,⁶ Stephen M. Rosenthal,⁷ Joshua D. Safer,⁸ Vin Tangpricha,⁹ and Guy G. T'Sjoen¹⁰

¹New York Presbyterian Hospital, Columbia University Medical Center, New York, New York 10032 (Retired); ²VU University Medical Center, 1007 MB Amsterdam, Netherlands (Retired); ³VU University Medical Center, 1007 MB Amsterdam, Netherlands (Retired); ⁴Leiden University Medical Center, 2300 RC Leiden, Netherlands; ⁵University of Texas Medical Branch, Galveston, Texas 77555; ⁶Mayo Clinic Evidence-Based Practice Center, Rochester, Minnesota 55905; ⁷University of California San Francisco, Benioff Children's Hospital, San Francisco, California 94143; ⁸Boston University School of Medicine, Boston, Massachusetts 02118; ⁹Emory University School of Medicine and the Atlanta VA Medical Center, Atlanta, Georgia 30322; and ¹⁰Ghent University Hospital, 9000 Ghent, Belgium

***Cosponsoring Associations:** American Association of Clinical Endocrinologists, American Society of Andrology, European Society for Pediatric Endocrinology, European Society of Endocrinology, Pediatric Endocrine Society, and World Professional Association for Transgender Health.

Objective: To update the "Endocrine Treatment of Transsexual Persons: An Endocrine Society Clinical Practice Guideline," published by the Endocrine Society in 2009.

Participants: The participants include an Endocrine Society–appointed task force of nine experts, a methodologist, and a medical writer.

Evidence: This evidence-based guideline was developed using the Grading of Recommendations, Assessment, Development, and Evaluation approach to describe the strength of recommendations and the quality of evidence. The task force commissioned two systematic reviews and used the best available evidence from other published systematic reviews and individual studies.

Consensus Process: Group meetings, conference calls, and e-mail communications enabled consensus. Endocrine Society committees, members and cosponsoring organizations reviewed and commented on preliminary drafts of the guidelines.

Conclusion: Gender affirmation is multidisciplinary treatment in which endocrinologists play an important role. Gender-dysphoric/gender-incongruent persons seek and/or are referred to endocrinologists to develop the physical characteristics of the affirmed gender. They require a safe and effective hormone regimen that will (1) suppress endogenous sex hormone secretion determined by the person's genetic/gonadal sex and (2) maintain sex hormone levels within the normal range for the person's affirmed gender. Hormone treatment is not recommended for prepubertal gender-dysphoric/gender-incongruent persons. Those clinicians who recommend gender-affirming endocrine treatments—appropriately trained diagnosing clinicians (required), a mental health provider for adolescents (required) and mental health

professional for adults (recommended)—should be knowledgeable about the diagnostic criteria and criteria for gender-affirming treatment, have sufficient training and experience in assessing psychopathology, and be willing to participate in the ongoing care throughout the endocrine transition. We recommend treating gender-dysphoric/gender-incongruent adolescents who have entered puberty at Tanner Stage G2/B2 by suppression with gonadotropin-releasing hormone agonists. Clinicians may add gender-affirming hormones after a multidisciplinary team has confirmed the persistence of gender dysphoria/gender incongruence and sufficient mental capacity to give informed consent to this partially irreversible treatment. Most adolescents have this capacity by age 16 years old. We recognize that there may be compelling reasons to initiate sex hormone treatment prior to age 16 years, although there is minimal published experience treating prior to 13.5 to 14 years of age. For the care of peripubertal youths and older adolescents, we recommend that an expert multidisciplinary team comprised of medical professionals and mental health professionals manage this treatment. The treating physician must confirm the criteria for treatment used by the referring mental health practitioner and collaborate with them in decisions about gender-affirming surgery in older adolescents. For adult gender-dysphoric/gender-incongruent persons, the treating clinicians (collectively) should have expertise in transgender-specific diagnostic criteria, mental health, primary care, hormone treatment, and surgery, as needed by the patient. We suggest maintaining physiologic levels of gender-appropriate hormones and monitoring for known risks and complications. When high doses of sex steroids are required to suppress endogenous sex steroids and/or in advanced age, clinicians may consider surgically removing natal gonads along with reducing sex steroid treatment. Clinicians should monitor both transgender males (female to male) and transgender females (male to female) for reproductive organ cancer risk when surgical removal is incomplete. Additionally, clinicians should persistently monitor adverse effects of sex steroids. For gender-affirming surgeries in adults, the treating physician must collaborate with and confirm the criteria for treatment used by the referring physician. Clinicians should avoid harming individuals (via hormone treatment) who have conditions other than gender dysphoria/gender incongruence and who may not benefit from the physical changes associated with this treatment. (*J Clin Endocrinol Metab* 102: 3869–3903, 2017)

Summary of Recommendations

1.0 Evaluation of youth and adults

- 1.1. We advise that only trained mental health professionals (MHPs) who meet the following criteria should diagnose gender dysphoria (GD)/gender incongruence in adults: (1) competence in using the Diagnostic and Statistical Manual of Mental Disorders (DSM) and/or the International Statistical Classification of Diseases and Related Health Problems (ICD) for diagnostic purposes, (2) the ability to diagnose GD/gender incongruence and make a distinction between GD/gender incongruence and conditions that have similar features (*e.g.*, body dysmorphic disorder), (3) training in diagnosing psychiatric conditions, (4) the ability to undertake or refer for appropriate treatment, (5) the ability to psychosocially assess the person's understanding, mental health, and social conditions that can impact gender-affirming hormone therapy, and (6) a practice of regularly attending relevant professional meetings. (Ungraded Good Practice Statement)
- 1.2. We advise that only MHPs who meet the following criteria should diagnose GD/gender incongruence in children and adolescents: (1) training in child and adolescent developmental psychology and psychopathology, (2) competence in using the DSM and/or the ICD for diagnostic purposes, (3) the ability to make a distinction between GD/gender incongruence and conditions that have similar features (*e.g.*, body dysmorphic disorder), (4) training in diagnosing psychiatric conditions, (5) the ability to undertake or refer for appropriate treatment, (6) the ability to psychosocially assess the person's understanding and social conditions that can impact gender-affirming hormone therapy, (7) a practice of regularly attending relevant professional meetings, and (8) knowledge of the criteria for puberty blocking and gender-affirming hormone treatment in adolescents. (Ungraded Good Practice Statement)
- 1.3. We advise that decisions regarding the social transition of prepubertal youths with GD/gender incongruence are made with the assistance of an MHP or another experienced professional. (Ungraded Good Practice Statement).

- 1.4. We recommend against puberty blocking and gender-affirming hormone treatment in pre-pubertal children with GD/gender incongruence. (1 ⊕⊕○○)
- 1.5. We recommend that clinicians inform and counsel all individuals seeking gender-affirming medical treatment regarding options for fertility preservation prior to initiating puberty suppression in adolescents and prior to treating with hormonal therapy of the affirmed gender in both adolescents and adults. (1 ⊕⊕⊕○)

2.0 Treatment of adolescents

- 2.1. We suggest that adolescents who meet diagnostic criteria for GD/gender incongruence, fulfill criteria for treatment, and are requesting treatment should initially undergo treatment to suppress pubertal development. (2 ⊕⊕○○)
- 2.2. We suggest that clinicians begin pubertal hormone suppression after girls and boys first exhibit physical changes of puberty. (2 ⊕⊕○○)
- 2.3. We recommend that, where indicated, GnRH analogues are used to suppress pubertal hormones. (1 ⊕⊕○○)
- 2.4. In adolescents who request sex hormone treatment (given this is a partly irreversible treatment), we recommend initiating treatment using a gradually increasing dose schedule after a multidisciplinary team of medical and MHPs has confirmed the persistence of GD/gender incongruence and sufficient mental capacity to give informed consent, which most adolescents have by age 16 years. (1 ⊕⊕○○).
- 2.5. We recognize that there may be compelling reasons to initiate sex hormone treatment prior to the age of 16 years in some adolescents with GD/gender incongruence, even though there are minimal published studies of gender-affirming hormone treatments administered before age 13.5 to 14 years. As with the care of adolescents ≥16 years of age, we recommend that an expert multidisciplinary team of medical and MHPs manage this treatment. (1 ⊕○○○)
- 2.6. We suggest monitoring clinical pubertal development every 3 to 6 months and laboratory parameters every 6 to 12 months during sex hormone treatment. (2 ⊕⊕○○)

3.0 Hormonal therapy for transgender adults

- 3.1. We recommend that clinicians confirm the diagnostic criteria of GD/gender incongruence and

- the criteria for the endocrine phase of gender transition before beginning treatment. (1 ⊕⊕⊕○)
- 3.2. We recommend that clinicians evaluate and address medical conditions that can be exacerbated by hormone depletion and treatment with sex hormones of the affirmed gender before beginning treatment. (1 ⊕⊕⊕○)
- 3.3. We suggest that clinicians measure hormone levels during treatment to ensure that endogenous sex steroids are suppressed and administered sex steroids are maintained in the normal physiologic range for the affirmed gender. (2 ⊕⊕○○)
- 3.4. We suggest that endocrinologists provide education to transgender individuals undergoing treatment about the onset and time course of physical changes induced by sex hormone treatment. (2 ⊕○○○)

4.0 Adverse outcome prevention and long-term care

- 4.1. We suggest regular clinical evaluation for physical changes and potential adverse changes in response to sex steroid hormones and laboratory monitoring of sex steroid hormone levels every 3 months during the first year of hormone therapy for transgender males and females and then once or twice yearly. (2 ⊕⊕○○)
- 4.2. We suggest periodically monitoring prolactin levels in transgender females treated with estrogens. (2 ⊕⊕○○)
- 4.3. We suggest that clinicians evaluate transgender persons treated with hormones for cardiovascular risk factors using fasting lipid profiles, diabetes screening, and/or other diagnostic tools. (2 ⊕⊕○○)
- 4.4. We recommend that clinicians obtain bone mineral density (BMD) measurements when risk factors for osteoporosis exist, specifically in those who stop sex hormone therapy after gonadectomy. (1 ⊕⊕○○)
- 4.5. We suggest that transgender females with no known increased risk of breast cancer follow breast-screening guidelines recommended for non-transgender females. (2 ⊕⊕○○)
- 4.6. We suggest that transgender females treated with estrogens follow individualized screening according to personal risk for prostatic disease and prostate cancer. (2 ⊕○○○)
- 4.7. We advise that clinicians determine the medical necessity of including a total hysterectomy and oophorectomy as part of gender-affirming surgery. (Ungraded Good Practice Statement)

5.0 Surgery for sex reassignment and gender confirmation

- 5.1. We recommend that a patient pursue genital gender-affirming surgery only after the MHP and the clinician responsible for endocrine transition therapy both agree that surgery is medically necessary and would benefit the patient's overall health and/or well-being. (1 ⊕⊕○○)
- 5.2. We advise that clinicians approve genital gender-affirming surgery only after completion of at least 1 year of consistent and compliant hormone treatment, unless hormone therapy is not desired or medically contraindicated. (Ungraded Good Practice Statement)
- 5.3. We advise that the clinician responsible for endocrine treatment and the primary care provider ensure appropriate medical clearance of transgender individuals for genital gender-affirming surgery and collaborate with the surgeon regarding hormone use during and after surgery. (Ungraded Good Practice Statement)
- 5.4. We recommend that clinicians refer hormone-treated transgender individuals for genital surgery when: (1) the individual has had a satisfactory social role change, (2) the individual is satisfied about the hormonal effects, and (3) the individual desires definitive surgical changes. (1 ⊕○○○)
- 5.5. We suggest that clinicians delay gender-affirming genital surgery involving gonadectomy and/or hysterectomy until the patient is at least 18 years old or legal age of majority in his or her country. (2 ⊕⊕○○)
- 5.6. We suggest that clinicians determine the timing of breast surgery for transgender males based upon the physical and mental health status of the individual. There is insufficient evidence to recommend a specific age requirement. (2 ⊕○○○)

Changes Since the Previous Guideline

Both the current guideline and the one published in 2009 contain similar sections. Listed here are the sections contained in the current guideline and the corresponding number of recommendations: Introduction, Evaluation of Youth and Adults (5), Treatment of Adolescents (6), Hormonal Therapy for Transgender Adults (4), Adverse Outcomes Prevention and Long-term Care (7), and Surgery for Sex Reassignment and Gender Confirmation (6). The current introduction updates the diagnostic classification of "gender dysphoria/gender incongruence." It also reviews the development of "gender identity" and summarizes its natural development. The section on

clinical evaluation of both youth and adults, defines in detail the professional qualifications required of those who diagnose and treat both adolescents and adults. We advise that decisions regarding the social transition of prepubertal youth are made with the assistance of a mental health professional or similarly experienced professional. We recommend against puberty blocking followed by gender-affirming hormone treatment of prepubertal children. Clinicians should inform pubertal children, adolescents, and adults seeking gender-confirming treatment of their options for fertility preservation. Prior to treatment, clinicians should evaluate the presence of medical conditions that may be worsened by hormone depletion and/or treatment. A multidisciplinary team, preferably composed of medical and mental health professionals, should monitor treatments. Clinicians evaluating transgender adults for endocrine treatment should confirm the diagnosis of persistent gender dysphoria/gender incongruence. Physicians should educate transgender persons regarding the time course of steroid-induced physical changes. Treatment should include periodic monitoring of hormone levels and metabolic parameters, as well as assessments of bone density and the impact upon prostate, gonads, and uterus. We also make recommendations for transgender persons who plan genital gender-affirming surgery.

Method of Development of Evidence-Based Clinical Practice Guidelines

The Clinical Guidelines Subcommittee (CGS) of the Endocrine Society deemed the diagnosis and treatment of individuals with GD/gender incongruence a priority area for revision and appointed a task force to formulate evidence based recommendations. The task force followed the approach recommended by the Grading of Recommendations, Assessment, Development, and Evaluation group, an international group with expertise in the development and implementation of evidence based guidelines (1). A detailed description of the grading scheme has been published elsewhere (2). The task force used the best available research evidence to develop the recommendations. The task force also used consistent language and graphical descriptions of both the strength of a recommendation and the quality of evidence. In terms of the strength of the recommendation, strong recommendations use the phrase "we recommend" and the number 1, and weak recommendations use the phrase "we suggest" and the number 2. Cross filled circles indicate the quality of the evidence, such that ⊕○○○ denotes very low quality evidence; ⊕⊕○○, low quality; ⊕⊕⊕○, moderate quality; and ⊕⊕⊕⊕, high quality. The task force has confidence that persons who receive care according to the strong recommendations will derive, on average, more benefit than harm. Weak recommendations require more careful consideration of the person's circumstances, values, and preferences to determine the best course of action. Linked to each recommendation is a description of the evidence and the

values that the task force considered in making the recommendation. In some instances, there are remarks in which the task force offers technical suggestions for testing conditions, dosing, and monitoring. These technical comments reflect the best available evidence applied to a typical person being treated. Often this evidence comes from the unsystematic observations of the task force and their preferences; therefore, one should consider these remarks as suggestions.

In this guideline, the task force made several statements to emphasize the importance of shared decision making, general preventive care measures, and basic principles of the treatment of transgender persons. They labeled these “Ungraded Good Practice Statement.” Direct evidence for these statements was either unavailable or not systematically appraised and considered out of the scope of this guideline. The intention of these statements is to draw attention to these principles.

The Endocrine Society maintains a rigorous conflict of interest review process for developing clinical practice guidelines. All task force members must declare any potential conflicts of interest by completing a conflict of interest form. The CGS reviews all conflicts of interest before the Society’s Council approves the members to participate on the task force and periodically during the development of the guideline. All others participating in the guideline’s development must also disclose any conflicts of interest in the matter under study, and most of these participants must be without any conflicts of interest. The CGS and the task force have reviewed all disclosures for this guideline and resolved or managed all identified conflicts of interest.

Conflicts of interest are defined as remuneration in any amount from commercial interests; grants; research support; consulting fees; salary; ownership interests [e.g., stocks and stock options (excluding diversified mutual funds)]; honoraria and other payments for participation in speakers’ bureaus, advisory boards, or boards of directors; and all other financial benefits. Completed forms are available through the Endocrine Society office.

The Endocrine Society provided the funding for this guideline; the task force received no funding or remuneration from commercial or other entities.

Commissioned Systematic Review

The task force commissioned two systematic reviews to support this guideline. The first one aimed to summarize the available evidence on the effect of sex steroid use in transgender individuals on lipids and cardiovascular outcomes. The review identified 29 eligible studies at moderate risk of bias. In transgender males (female to male), sex steroid therapy was associated with a statistically significant increase in serum triglycerides and low-density lipoprotein cholesterol levels. High-density lipoprotein cholesterol levels decreased significantly across all follow-up time periods. In transgender females (male to female), serum triglycerides were significantly higher without any changes in other parameters. Few myocardial infarction, stroke, venous thromboembolism (VTE), and death events were reported. These events were more frequent in transgender females. However, the

quality of the evidence was low. The second review summarized the available evidence regarding the effect of sex steroids on bone health in transgender individuals and identified 13 studies. In transgender males, there was no statistically significant difference in the lumbar spine, femoral neck, or total hip BMD at 12 and 24 months compared with baseline values before initiating masculinizing hormone therapy. In transgender females, there was a statistically significant increase in lumbar spine BMD at 12 months and 24 months compared with baseline values before initiation of feminizing hormone therapy. There was minimal information on fracture rates. The quality of evidence was also low.

Introduction

Throughout recorded history (in the absence of an endocrine disorder) some men and women have experienced confusion and anguish resulting from rigid, forced conformity to sexual dimorphism. In modern history, there have been numerous ongoing biological, psychological, cultural, political, and sociological debates over various aspects of gender variance. The 20th century marked the emergence of a social awakening for men and women with the belief that they are “trapped” in the wrong body (3). Magnus Hirschfeld and Harry Benjamin, among others, pioneered the medical responses to those who sought relief from and a resolution to their profound discomfort. Although the term transsexual became widely known after Benjamin wrote “The Transsexual Phenomenon” (4), it was Hirschfeld who coined the term “transsexual” in 1923 to describe people who want to live a life that corresponds with their experienced gender vs their designated gender (5). Magnus Hirschfeld (6) and others (4, 7) have described other types of trans phenomena besides transsexualism. These early researchers proposed that the gender identity of these people was located somewhere along a unidimensional continuum. This continuum ranged from all male through “something in between” to all female. Yet such a classification does not take into account that people may have gender identities outside this continuum. For instance, some experience themselves as having both a male and female gender identity, whereas others completely renounce any gender classification (8, 9). There are also reports of individuals experiencing a continuous and rapid involuntary alternation between a male and female identity (10) or men who do not experience themselves as men but do not want to live as women (11, 12). In some countries, (e.g., Nepal, Bangladesh, and Australia), these nonmale or nonfemale genders are officially recognized (13). Specific treatment protocols, however, have not yet been developed for these groups.

Instead of the term transsexualism, the current classification system of the American Psychiatric Association uses the term gender dysphoria in its diagnosis of persons who are not satisfied with their designated gender (14). The current version of the World Health Organization's ICD-10 still uses the term transsexualism when diagnosing adolescents and adults. However, for the ICD-11, the World Health Organization has proposed using the term "gender incongruence" (15).

Treating persons with GD/gender incongruence (15) was previously limited to relatively ineffective elixirs or creams. However, more effective endocrinology-based treatments became possible with the availability of testosterone in 1935 and diethylstilbestrol in 1938. Reports of individuals with GD/gender incongruence who were treated with hormones and gender-affirming surgery appeared in the press during the second half of the 20th century. The Harry Benjamin International Gender Dysphoria Association was founded in September 1979 and is now called the World Professional Association for Transgender Health (WPATH). WPATH published its first Standards of Care in 1979. These standards have since been regularly updated, providing guidance for treating persons with GD/gender incongruence (16).

Prior to 1975, few peer-reviewed articles were published concerning endocrine treatment of transgender persons. Since then, more than two thousand articles about various aspects of transgender care have appeared.

It is the purpose of this guideline to make detailed recommendations and suggestions, based on existing medical literature and clinical experience, that will enable treating physicians to maximize benefit and minimize risk when caring for individuals diagnosed with GD/gender incongruence.

In the future, we need more rigorous evaluations of the effectiveness and safety of endocrine and surgical protocols. Specifically, endocrine treatment protocols for GD/gender incongruence should include the careful assessment of the following: (1) the effects of prolonged delay of puberty in adolescents on bone health, gonadal function, and the brain (including effects on cognitive, emotional, social, and sexual development); (2) the effects of treatment in adults on sex hormone levels; (3) the requirement for and the effects of progestins and other agents used to suppress endogenous sex steroids during treatment; and (4) the risks and benefits of gender-affirming hormone treatment in older transgender people.

To successfully establish and enact these protocols, a commitment of mental health and endocrine investigators is required to collaborate in long-term, large-scale

studies across countries that use the same diagnostic and inclusion criteria, medications, assay methods, and response assessment tools (*e.g.*, the European Network for the Investigation of Gender Incongruence) (17, 18).

Terminology and its use vary and continue to evolve. Table 1 contains the definitions of terms as they are used throughout this guideline.

Biological Determinants of Gender Identity Development

One's self-awareness as male or female changes gradually during infant life and childhood. This process of cognitive and affective learning evolves with interactions with parents, peers, and environment. A fairly accurate timetable exists outlining the steps in this process (19). Normative psychological literature, however, does not address if and when gender identity becomes crystallized and what factors contribute to the development of a gender identity that is not congruent with the gender of rearing. Results of studies from a variety of biomedical disciplines—genetic, endocrine, and neuroanatomic—support the concept that gender identity and/or gender expression (20) likely reflect a complex interplay of biological, environmental, and cultural factors (21, 22).

With respect to endocrine considerations, studies have failed to find differences in circulating levels of sex steroids between transgender and nontransgender individuals (23). However, studies in individuals with a disorder/difference of sex development (DSD) have informed our understanding of the role that hormones may play in gender identity outcome, even though most persons with GD/gender incongruence do not have a DSD. For example, although most 46,XX adult individuals with virilizing congenital adrenal hyperplasia caused by mutations in *CYP21A2* reported a female gender identity, the prevalence of GD/gender incongruence was much greater in this group than in the general population without a DSD. This supports the concept that there is a role for prenatal/postnatal androgens in gender development (24–26), although some studies indicate that prenatal androgens are more likely to affect gender behavior and sexual orientation rather than gender identity *per se* (27, 28).

Researchers have made similar observations regarding the potential role of androgens in the development of gender identity in other individuals with DSD. For example, a review of two groups of 46,XY persons, each with androgen synthesis deficiencies and female raised, reported transgender male (female-to-male) gender role changes in 56% to 63% and 39% to 64% of patients, respectively (29). Also, in 46,XY female-raised individuals with cloacal

Table 1. Definitions of Terms Used in This Guideline

Biological sex, biological male or female: These terms refer to physical aspects of maleness and femaleness. As these may not be in line with each other (e.g., a person with XY chromosomes may have female-appearing genitalia), the terms biological sex and biological male or female are imprecise and should be avoided.

Cisgender: This means not transgender. An alternative way to describe individuals who are not transgender is “non-transgender people.”

Gender-affirming (hormone) treatment: See “gender reassignment”

Gender dysphoria: This is the distress and unease experienced if gender identity and designated gender are not completely congruent (see Table 2). In 2013, the American Psychiatric Association released the fifth edition of the DSM-5, which replaced “gender identity disorder” with “gender dysphoria” and changed the criteria for diagnosis.

Gender expression: This refers to external manifestations of gender, expressed through one’s name, pronouns, clothing, haircut, behavior, voice, or body characteristics. Typically, transgender people seek to make their gender expression align with their gender identity, rather than their designated gender.

Gender identity/experienced gender: This refers to one’s internal, deeply held sense of gender. For transgender people, their gender identity does not match their sex designated at birth. Most people have a gender identity of man or woman (or boy or girl). For some people, their gender identity does not fit neatly into one of those two choices. Unlike gender expression (see below), gender identity is not visible to others.

Gender identity disorder: This is the term used for GD/gender incongruence in previous versions of DSM (see “gender dysphoria”). The ICD-10 still uses the term for diagnosing child diagnoses, but the upcoming ICD-11 has proposed using “gender incongruence of childhood.”

Gender incongruence: This is an umbrella term used when the gender identity and/or gender expression differs from what is typically associated with the designated gender. Gender incongruence is also the proposed name of the gender identity–related diagnoses in ICD-11. Not all individuals with gender incongruence have gender dysphoria or seek treatment.

Gender variance: See “gender incongruence”

Gender reassignment: This refers to the treatment procedure for those who want to adapt their bodies to the experienced gender by means of hormones and/or surgery. This is also called gender-confirming or gender-affirming treatment.

Gender-reassignment surgery (gender-confirming/gender-affirming surgery): These terms refer only to the surgical part of gender-confirming/gender-affirming treatment.

Gender role: This refers to behaviors, attitudes, and personality traits that a society (in a given culture and historical period) designates as masculine or feminine and/or that society associates with or considers typical of the social role of men or women.

Sex designated at birth: This refers to sex assigned at birth, usually based on genital anatomy.

Sex: This refers to attributes that characterize biological maleness or femaleness. The best known attributes include the sex-determining genes, the sex chromosomes, the H-Y antigen, the gonads, sex hormones, internal and external genitalia, and secondary sex characteristics.

Sexual orientation: This term describes an individual’s enduring physical and emotional attraction to another person. Gender identity and sexual orientation are not the same. Irrespective of their gender identity, transgender people may be attracted to women (gynephilic), attracted to men (androphilic), bisexual, asexual, or queer.

Transgender: This is an umbrella term for people whose gender identity and/or gender expression differs from what is typically associated with their sex designated at birth. Not all transgender individuals seek treatment.

Transgender male (also: trans man, female-to-male, transgender male): This refers to individuals assigned female at birth but who identify and live as men.

Transgender woman (also: trans woman, male-to-female, transgender female): This refers to individuals assigned male at birth but who identify and live as women.

Transition: This refers to the process during which transgender persons change their physical, social, and/or legal characteristics consistent with the affirmed gender identity. Prepubertal children may choose to transition socially.

Transsexual: This is an older term that originated in the medical and psychological communities to refer to individuals who have permanently transitioned through medical interventions or desired to do so.

exstrophy and penile agenesis, the occurrence of transgender male changes was significantly more prevalent than in the general population (30, 31). However, the fact that a high percentage of individuals with the same conditions did not change gender suggests that cultural factors may play a role as well.

With respect to genetics and gender identity, several studies have suggested heritability of GD/gender incongruence (32, 33). In particular, a study by Heylens *et al.* (33) demonstrated a 39.1% concordance rate for gender identity disorder (based on the DSM-IV criteria) in 23 monozygotic twin pairs but no concordance in 21 same-sex dizygotic or seven opposite-sex twin pairs. Although numerous investigators have sought to identify

specific genes associated with GD/gender incongruence, such studies have been inconsistent and without strong statistical significance (34–38).

Studies focusing on brain structure suggest that the brain phenotypes of people with GD/gender incongruence differ in various ways from control males and females, but that there is not a complete sex reversal in brain structures (39).

In summary, although there is much that is still unknown with respect to gender identity and its expression, compelling studies support the concept that biologic factors, in addition to environmental factors, contribute to this fundamental aspect of human development.

Natural History of Children With GD/Gender Incongruence

With current knowledge, we cannot predict the psychosexual outcome for any specific child. Prospective follow-up studies show that childhood GD/gender incongruence does not invariably persist into adolescence and adulthood (so-called “desisters”). Combining all outcome studies to date, the GD/gender incongruence of a minority of prepubertal children appears to persist in adolescence (20, 40). In adolescence, a significant number of these desisters identify as homosexual or bisexual. It may be that children who only showed some gender nonconforming characteristics have been included in the follow-up studies, because the DSM-IV text revision criteria for a diagnosis were rather broad. However, the persistence of GD/gender incongruence into adolescence is more likely if it had been extreme in childhood (41, 42). With the newer, stricter criteria of the DSM-5 (Table 2), persistence rates may well be different in future studies.

1.0 Evaluation of Youth and Adults

Gender-affirming treatment is a multidisciplinary effort. After evaluation, education, and diagnosis, treatment may include mental health care, hormone therapy, and/or surgical therapy. Together with an MHP, hormone-prescribing clinicians should examine the psychosocial impact of the potential changes on people’s lives, including mental health, friends, family, jobs, and their role in society. Transgender individuals should be encouraged to experience living in the new gender role and assess whether

this improves their quality of life. Although the focus of this guideline is gender-affirming hormone therapy, collaboration with appropriate professionals responsible for each aspect of treatment maximizes a successful outcome.

Diagnostic assessment and mental health care

GD/gender incongruence may be accompanied with psychological or psychiatric problems (43–51). It is therefore necessary that clinicians who prescribe hormones and are involved in diagnosis and psychosocial assessment meet the following criteria: (1) are competent in using the DSM and/or the ICD for diagnostic purposes, (2) are able to diagnose GD/gender incongruence and make a distinction between GD/gender incongruence and conditions that have similar features (*e.g.*, body dysmorphic disorder), (3) are trained in diagnosing psychiatric conditions, (4) undertake or refer for appropriate treatment, (5) are able to do a psychosocial assessment of the patient’s understanding, mental health, and social conditions that can impact gender-affirming hormone therapy, and (6) regularly attend relevant professional meetings.

Because of the psychological vulnerability of many individuals with GD/gender incongruence, it is important that mental health care is available before, during, and sometimes also after transitioning. For children and adolescents, an MHP who has training/experience in child and adolescent gender development (as well as child and adolescent psychopathology) should make the diagnosis, because assessing GD/gender incongruence in children and adolescents is often extremely complex.

During assessment, the clinician obtains information from the individual seeking gender-affirming treatment. In the case

Table 2. DSM-5 Criteria for Gender Dysphoria in Adolescents and Adults

-
- A. A marked incongruence between one’s experienced/expressed gender and natal gender of at least 6 mo in duration, as manifested by at least two of the following:
1. A marked incongruence between one’s experienced/expressed gender and primary and/or secondary sex characteristics (or in young adolescents, the anticipated secondary sex characteristics)
 2. A strong desire to be rid of one’s primary and/or secondary sex characteristics because of a marked incongruence with one’s experienced/expressed gender (or in young adolescents, a desire to prevent the development of the anticipated secondary sex characteristics)
 3. A strong desire for the primary and/or secondary sex characteristics of the other gender
 4. A strong desire to be of the other gender (or some alternative gender different from one’s designated gender)
 5. A strong desire to be treated as the other gender (or some alternative gender different from one’s designated gender)
 6. A strong conviction that one has the typical feelings and reactions of the other gender (or some alternative gender different from one’s designated gender)
- B. The condition is associated with clinically significant distress or impairment in social, occupational, or other important areas of functioning.
- Specify if:
1. The condition exists with a disorder of sex development.
 2. The condition is posttransitional, in that the individual has transitioned to full-time living in the desired gender (with or without legalization of gender change) and has undergone (or is preparing to have) at least one sex-related medical procedure or treatment regimen—namely, regular sex hormone treatment or gender reassignment surgery confirming the desired gender (*e.g.*, penectomy, vaginoplasty in natal males; mastectomy or phalloplasty in natal females).
-

of adolescents, the clinician also obtains information from the parents or guardians regarding various aspects of the child's general and psychosexual development and current functioning. On the basis of this information, the clinician:

- decides whether the individual fulfills criteria for treatment (see Tables 2 and 3) for GD/gender incongruence (DSM-5) or transsexualism (DSM-5 and/or ICD-10);
- informs the individual about the possibilities and limitations of various kinds of treatment (hormonal/surgical and nonhormonal), and if medical treatment is desired, provides correct information to prevent unrealistically high expectations;
- assesses whether medical interventions may result in unfavorable psychological and social outcomes.

In cases in which severe psychopathology, circumstances, or both seriously interfere with the diagnostic work or make satisfactory treatment unlikely, clinicians should assist the adolescent in managing these other issues. Literature on postoperative regret suggests that besides poor quality of surgery, severe psychiatric comorbidity and lack of support may interfere with positive outcomes (52–56).

For adolescents, the diagnostic procedure usually includes a complete psychodiagnostic assessment (57) and an assessment of the decision-making capability of the youth. An evaluation to assess the family's ability to endure stress, give support, and deal with the complexities of the adolescent's situation should be part of the diagnostic phase (58).

Social transitioning

A change in gender expression and role (which may involve living part time or full time in another gender role that is consistent with one's gender identity) may test the person's resolve, the capacity to function in the affirmed gender, and the adequacy of social, economic, and psychological supports. It assists both the individual and the clinician in their judgments about how to proceed (16). During social transitioning, the person's feelings about the social transformation (including coping with the responses of others) is a major focus of the counseling. The optimal timing for social transitioning may differ between individuals. Sometimes people wait until they

start gender-affirming hormone treatment to make social transitioning easier, but individuals increasingly start social transitioning long before they receive medically supervised, gender-affirming hormone treatment.

Criteria

Adolescents and adults seeking gender-affirming hormone treatment and surgery should satisfy certain criteria before proceeding (16). Criteria for gender-affirming hormone therapy for adults are in Table 4, and criteria for gender-affirming hormone therapy for adolescents are in Table 5. Follow-up studies in adults meeting these criteria indicate a high satisfaction rate with treatment (59). However, the quality of evidence is usually low. A few follow-up studies on adolescents who fulfilled these criteria also indicated good treatment results (60–63).

Recommendations for Those Involved in the Gender-Affirming Hormone Treatment of Individuals With GD/Gender Incongruence

- 1.1. We advise that only trained MHPs who meet the following criteria should diagnose GD/gender incongruence in adults: (1) competence in using the DSM and/or the ICD for diagnostic purposes, (2) the ability to diagnose GD/gender incongruence and make a distinction between GD/gender incongruence and conditions that have similar features (*e.g.*, body dysmorphic disorder), (3) training in diagnosing psychiatric conditions, (4) the ability to undertake or refer for appropriate treatment, (5) the ability to psychosocially assess the person's understanding, mental health, and social conditions that can impact gender-affirming hormone therapy, and (6) a practice of regularly attending relevant professional meetings. (Ungraded Good Practice Statement)
- 1.2. We advise that only MHPs who meet the following criteria should diagnose GD/gender incongruence in children and adolescents: (1) training in child and adolescent developmental psychology and psychopathology, (2) competence in using the DSM and/or ICD for diagnostic

Table 3. ICD-10 Criteria for Transsexualism

Transsexualism (F64.0) has three criteria:

1. The desire to live and be accepted as a member of the opposite sex, usually accompanied by the wish to make his or her body as congruent as possible with the preferred sex through surgery and hormone treatments.
2. The transsexual identity has been present persistently for at least 2 y.
3. The disorder is not a symptom of another mental disorder or a genetic, DSD, or chromosomal abnormality.

Table 4. Criteria for Gender-Affirming Hormone Therapy for Adults

1. Persistent, well-documented gender dysphoria/gender incongruence
2. The capacity to make a fully informed decision and to consent for treatment
3. The age of majority in a given country (if younger, follow the criteria for adolescents)
4. Mental health concerns, if present, must be reasonably well controlled

Reproduced from World Professional Association for Transgender Health (16).

purposes, (3) the ability to make a distinction between GD/gender incongruence and conditions that have similar features (*e.g.*, body dysmorphic disorder), (4) training in diagnosing psychiatric conditions, (5) the ability to undertake or refer for appropriate treatment, (6) the ability to psychosocially assess the person's understanding and social conditions that can impact gender-affirming hormone therapy, (7) a practice of regularly attending relevant professional meetings, and (8) knowledge of the criteria for puberty blocking and gender-affirming hormone treatment in adolescents. (Ungraded Good Practice Statement)

Evidence

Individuals with gender identity issues may have psychological or psychiatric problems (43–48, 50, 51, 64, 65). It is therefore necessary that clinicians making the diagnosis are able to make a distinction between GD/gender incongruence and conditions that have similar features. Examples of conditions with similar features are body dysmorphic disorder, body identity integrity disorder (a condition in which individuals have a sense that their anatomical configuration as an able-bodied person is somehow wrong or inappropriate) (66), or certain forms of eunuchism (in which a person is preoccupied with or engages in castration and/or penectomy for

Table 5. Criteria for Gender-Affirming Hormone Therapy for Adolescents

Adolescents are eligible for GnRH agonist treatment if:

1. A qualified MHP has confirmed that:
 - the adolescent has demonstrated a long-lasting and intense pattern of gender nonconformity or gender dysphoria (whether suppressed or expressed),
 - gender dysphoria worsened with the onset of puberty,
 - any coexisting psychological, medical, or social problems that could interfere with treatment (*e.g.*, that may compromise treatment adherence) have been addressed, such that the adolescent's situation and functioning are stable enough to start treatment,
 - the adolescent has sufficient mental capacity to give informed consent to this (reversible) treatment,
2. And the adolescent:
 - has been informed of the effects and side effects of treatment (including potential loss of fertility if the individual subsequently continues with sex hormone treatment) and options to preserve fertility,
 - has given informed consent and (particularly when the adolescent has not reached the age of legal medical consent, depending on applicable legislation) the parents or other caretakers or guardians have consented to the treatment and are involved in supporting the adolescent throughout the treatment process,
3. And a pediatric endocrinologist or other clinician experienced in pubertal assessment
 - agrees with the indication for GnRH agonist treatment,
 - has confirmed that puberty has started in the adolescent (Tanner stage \geq G2/B2),
 - has confirmed that there are no medical contraindications to GnRH agonist treatment.

Adolescents are eligible for subsequent sex hormone treatment if:

1. A qualified MHP has confirmed:
 - the persistence of gender dysphoria,
 - any coexisting psychological, medical, or social problems that could interfere with treatment (*e.g.*, that may compromise treatment adherence) have been addressed, such that the adolescent's situation and functioning are stable enough to start sex hormone treatment,
 - the adolescent has sufficient mental capacity (which most adolescents have by age 16 years) to estimate the consequences of this (partly) irreversible treatment, weigh the benefits and risks, and give informed consent to this (partly) irreversible treatment,
2. And the adolescent:
 - has been informed of the (irreversible) effects and side effects of treatment (including potential loss of fertility and options to preserve fertility),
 - has given informed consent and (particularly when the adolescent has not reached the age of legal medical consent, depending on applicable legislation) the parents or other caretakers or guardians have consented to the treatment and are involved in supporting the adolescent throughout the treatment process,
3. And a pediatric endocrinologist or other clinician experienced in pubertal induction:
 - agrees with the indication for sex hormone treatment,
 - has confirmed that there are no medical contraindications to sex hormone treatment.

Reproduced from World Professional Association for Transgender Health (16).

reasons that are not gender identity related) (11). Clinicians should also be able to diagnose psychiatric conditions accurately and ensure that these conditions are treated appropriately, particularly when the conditions may complicate treatment, affect the outcome of gender-affirming treatment, or be affected by hormone use.

Values and preferences

The task force placed a very high value on avoiding harm from hormone treatment in individuals who have conditions other than GD/gender incongruence and who may not benefit from the physical changes associated with this treatment and placed a low value on any potential benefit these persons believe they may derive from hormone treatment. This justifies the good practice statement.

- 1.3. We advise that decisions regarding the social transition of prepubertal youths with GD/gender incongruence are made with the assistance of an MHP or another experienced professional. (Ungraded Good Practice Statement).
- 1.4. We recommend against puberty blocking and gender-affirming hormone treatment in prepubertal children with GD/gender incongruence. (1 ⊕⊕○○)

Evidence

In most children diagnosed with GD/gender incongruence, it did not persist into adolescence. The percentages differed among studies, probably dependent on which version of the DSM clinicians used, the patient's age, the recruitment criteria, and perhaps cultural factors. However, the large majority (about 85%) of prepubertal children with a childhood diagnosis did not remain GD/gender incongruent in adolescence (20). If children have completely socially transitioned, they may have great difficulty in returning to the original gender role upon entering puberty (40). Social transition is associated with the persistence of GD/gender incongruence as a child progresses into adolescence. It may be that the presence of GD/gender incongruence in prepubertal children is the earliest sign that a child is destined to be transgender as an adolescent/adult (20). However, social transition (in addition to GD/gender incongruence) has been found to contribute to the likelihood of persistence.

This recommendation, however, does not imply that children should be discouraged from showing gender-variant behaviors or should be punished for exhibiting such behaviors. In individual cases, an early complete social transition may result in a more favorable outcome, but there are currently no criteria to identify the

GD/gender-incongruent children to whom this applies. At the present time, clinical experience suggests that persistence of GD/gender incongruence can only be reliably assessed after the first signs of puberty.

Values and preferences

The task force placed a high value on avoiding harm with gender-affirming hormone therapy in prepubertal children with GD/gender incongruence. This justifies the strong recommendation in the face of low-quality evidence.

- 1.5. We recommend that clinicians inform and counsel all individuals seeking gender-affirming medical treatment regarding options for fertility preservation prior to initiating puberty suppression in adolescents and prior to treating with hormonal therapy of the affirmed gender in both adolescents and adults. (1 ⊕⊕⊕○)

Remarks

Persons considering hormone use for gender affirmation need adequate information about this treatment in general and about fertility effects of hormone treatment in particular to make an informed and balanced decision (67, 68). Because young adolescents may not feel qualified to make decisions about fertility and may not fully understand the potential effects of hormonal interventions, consent and protocol education should include parents, the referring MHP(s), and other members of the adolescent's support group. To our knowledge, there are no formally evaluated decision aids available to assist in the discussion and decision regarding the future fertility of adolescents or adults beginning gender-affirming treatment.

Treating early pubertal youth with GnRH analogs will temporarily impair spermatogenesis and oocyte maturation. Given that an increasing number of transgender youth want to preserve fertility potential, delaying or temporarily discontinuing GnRH analogs to promote gamete maturation is an option. This option is often not preferred, because mature sperm production is associated with later stages of puberty and with the significant development of secondary sex characteristics.

For those designated male at birth with GD/gender incongruence and who are in early puberty, sperm production and the development of the reproductive tract are insufficient for the cryopreservation of sperm. However, prolonged pubertal suppression using GnRH analogs is reversible and clinicians should inform these individuals that sperm production can be initiated following prolonged gonadotropin suppression. This can be accomplished by spontaneous gonadotropin recovery after

cessation of GnRH analogs or by gonadotropin treatment and will probably be associated with physical manifestations of testosterone production, as stated above. Note that there are no data in this population concerning the time required for sufficient spermatogenesis to collect enough sperm for later fertility. In males treated for precocious puberty, spermarche was reported 0.7 to 3 years after cessation of GnRH analogs (69). In adult men with gonadotropin deficiency, sperm are noted in seminal fluid by 6 to 12 months of gonadotropin treatment. However, sperm numbers when partners of these patients conceive are far below the “normal range” (70, 71).

In girls, no studies have reported long-term, adverse effects of pubertal suppression on ovarian function after treatment cessation (72, 73). Clinicians should inform adolescents that no data are available regarding either time to spontaneous ovulation after cessation of GnRH analogs or the response to ovulation induction following prolonged gonadotropin suppression.

In males with GD/gender incongruence, when medical treatment is started in a later phase of puberty or in adulthood, spermatogenesis is sufficient for cryopreservation and storage of sperm. *In vitro* spermatogenesis is currently under investigation. Restoration of spermatogenesis after prolonged estrogen treatment has not been studied.

In females with GD/gender incongruence, the effect of prolonged treatment with exogenous testosterone on ovarian function is uncertain. There have been reports of an increased incidence of polycystic ovaries in transgender males, both prior to and as a result of androgen treatment (74–77), although these reports were not confirmed by others (78). Pregnancy has been reported in transgender males who have had prolonged androgen treatment and have discontinued testosterone but have not had genital surgery (79, 80). A reproductive endocrine gynecologist can counsel patients before gender-affirming hormone treatment or surgery regarding potential fertility options (81). Techniques for cryopreservation of oocytes, embryos, and ovarian tissue continue to improve, and oocyte maturation of immature tissue is being studied (82).

2.0 Treatment of Adolescents

During the past decade, clinicians have progressively acknowledged the suffering of young adolescents with GD/gender incongruence. In some forms of GD/gender incongruence, psychological interventions may be useful and sufficient. However, for many adolescents with GD/gender incongruence, the pubertal physical changes are unbearable. As early medical intervention may prevent

psychological harm, various clinics have decided to start treating young adolescents with GD/gender incongruence with puberty-suppressing medication (a GnRH analog). As compared with starting gender-affirming treatment long after the first phases of puberty, a benefit of pubertal suppression at early puberty may be a better psychological and physical outcome.

In girls, the first physical sign of puberty is the budding of the breasts followed by an increase in breast and fat tissue. Breast development is also associated with the pubertal growth spurt, and menarche occurs ~2 years later. In boys, the first physical change is testicular growth. A testicular volume ≥ 4 mL is seen as consistent with the initiation of physical puberty. At the beginning of puberty, estradiol and testosterone levels are still low and are best measured in the early morning with an ultrasensitive assay. From a testicular volume of 10 mL, daytime testosterone levels increase, leading to virilization (83). Note that pubic hair and/or axillary hair/odor may not reflect the onset of gonadarche; instead, it may reflect adrenarche alone.

- 2.1. We suggest that adolescents who meet diagnostic criteria for GD/gender incongruence, fulfill criteria for treatment (Table 5), and are requesting treatment should initially undergo treatment to suppress pubertal development. (2 ⊕⊕○○)
- 2.2. We suggest that clinicians begin pubertal hormone suppression after girls and boys first exhibit physical changes of puberty (Tanner stages G2/B2). (2 ⊕⊕○○)

Evidence

Pubertal suppression can expand the diagnostic phase by a long period, giving the subject more time to explore options and to live in the experienced gender before making a decision to proceed with gender-affirming sex hormone treatments and/or surgery, some of which is irreversible (84, 85). Pubertal suppression is fully reversible, enabling full pubertal development in the natal gender, after cessation of treatment, if appropriate. The experience of full endogenous puberty is an undesirable condition for the GD/gender-incongruent individual and may seriously interfere with healthy psychological functioning and well-being. Treating GD/gender-incongruent adolescents entering puberty with GnRH analogs has been shown to improve psychological functioning in several domains (86).

Another reason to start blocking pubertal hormones early in puberty is that the physical outcome is improved compared with initiating physical transition after puberty has been completed (60, 62). Looking like a man or woman when living as the opposite sex creates difficult

barriers with enormous life-long disadvantages. We therefore advise starting suppression in early puberty to prevent the irreversible development of undesirable secondary sex characteristics. However, adolescents with GD/gender incongruence should experience the first changes of their endogenous spontaneous puberty, because their emotional reaction to these first physical changes has diagnostic value in establishing the persistence of GD/gender incongruence (85). Thus, Tanner stage 2 is the optimal time to start pubertal suppression. However, pubertal suppression treatment in early puberty will limit the growth of the penis and scrotum, which will have a potential effect on future surgical treatments (87).

Clinicians can also use pubertal suppression in adolescents in later pubertal stages to stop menses in transgender males and prevent facial hair growth in transgender females. However, in contrast to the effects in early pubertal adolescents, physical sex characteristics (such as more advanced breast development in transgender boys and lowering of the voice and outgrowth of the jaw and brow in transgender girls) are not reversible.

Values and preferences

These recommendations place a high value on avoiding an unsatisfactory physical outcome when secondary sex characteristics have become manifest and irreversible, a higher value on psychological well-being, and a lower value on avoiding potential harm from early pubertal suppression.

Remarks

Table 6 lists the Tanner stages of breast and male genital development. Careful documentation of hallmarks of pubertal development will ensure precise timing when initiating pubertal suppression once puberty has started. Clinicians can use pubertal LH and sex steroid levels to confirm that puberty has progressed sufficiently before starting pubertal suppression (88). Reference

ranges for sex steroids by Tanner stage may vary depending on the assay used. Ultrasensitive sex steroid and gonadotropin assays will help clinicians document early pubertal changes.

Irreversible and, for GD/gender-incongruent adolescents, undesirable sex characteristics in female puberty are breasts, female body habitus, and, in some cases, relative short stature. In male puberty, they are a prominent Adam's apple; low voice; male bone configuration, such as a large jaw, big feet and hands, and tall stature; and male hair pattern on the face and extremities.

- 2.3. We recommend that, where indicated, GnRH analogues are used to suppress pubertal hormones. (1 ⊕⊕○○)

Evidence

Clinicians can suppress pubertal development and gonadal function most effectively via gonadotropin suppression using GnRH analogs. GnRH analogs are long-acting agonists that suppress gonadotropins by GnRH receptor desensitization after an initial increase of gonadotropins during ~10 days after the first and (to a lesser degree) the second injection (89). Antagonists immediately suppress pituitary gonadotropin secretion (90, 91). Long-acting GnRH analogs are the currently preferred treatment option. Clinicians may consider long-acting GnRH antagonists when evidence on their safety and efficacy in adolescents becomes available.

During GnRH analog treatment, slight development of secondary sex characteristics may regress, and in a later phase of pubertal development, it will stop. In girls, breast tissue will become atrophic, and menses will stop. In boys, virilization will stop, and testicular volume may decrease (92).

An advantage of using GnRH analogs is the reversibility of the intervention. If, after extensive exploration of his/her transition wish, the individual no longer desires transition, they can discontinue pubertal suppression. In subjects with

Table 6. Tanner Stages of Breast Development and Male External Genitalia

The description of Tanner stages for breast development:

1. Prepubertal
2. Breast and papilla elevated as small mound; areolar diameter increased
3. Breast and areola enlarged, no contour separation
4. Areola and papilla form secondary mound
5. Mature; nipple projects, areola part of general breast contour

For penis and testes:

1. Prepubertal, testicular volume <4 mL
2. Slight enlargement of penis; enlarged scrotum, pink, texture altered, testes 4–6 mL
3. Penis longer, testes larger (8–12 mL)
4. Penis and glans larger, including increase in breadth; testes larger (12–15 mL), scrotum dark
5. Penis adult size; testicular volume > 15 ml

Adapted from Lawrence (56).

precocious puberty, spontaneous pubertal development has been shown to resume after patients discontinue taking GnRH analogs (93).

Recommendations 2.1 to 2.3 are supported by a prospective follow-up study from The Netherlands. This report assessed mental health outcomes in 55 transgender adolescents/young adults (22 transgender females and 33 transgender males) at three time points: (1) before the start of GnRH agonist (average age of 14.8 years at start of treatment), (2) at initiation of gender-affirming hormones (average age of 16.7 years at start of treatment), and (3) 1 year after “gender-reassignment surgery” (average age of 20.7 years) (63). Despite a decrease in depression and an improvement in general mental health functioning, GD/gender incongruence persisted through pubertal suppression, as previously reported (86). However, following sex hormone treatment and gender-reassignment surgery, GD/gender incongruence was resolved and psychological functioning steadily improved (63). Furthermore, well-being was similar to or better than that reported by age-matched young adults from the general population, and none of the study participants regretted treatment. This study represents the first long-term follow-up of individuals managed according to currently existing clinical practice guidelines for transgender youth, and it underscores the benefit of the multidisciplinary approach pioneered in The Netherlands; however, further studies are needed.

Side effects

The primary risks of pubertal suppression in GD/gender-incongruent adolescents may include adverse effects on bone mineralization (which can theoretically be reversed with sex hormone treatment), compromised fertility if the person subsequently is treated with sex hormones, and unknown effects on brain development. Few data are available on the effect of GnRH analogs on BMD in adolescents with GD/gender incongruence. Initial data in GD/gender-incongruent subjects demonstrated no change of absolute areal BMD during 2 years of GnRH analog therapy but a decrease in BMD z scores (85). A recent study also suggested suboptimal bone mineral accrual during GnRH analog treatment. The study reported a decrease in areal BMD z scores and of bone mineral apparent density z scores (which takes the size of the bone into account) in 19 transgender males treated with GnRH analogs from a mean age of 15.0 years (standard deviation = 2.0 years) for a median duration of 1.5 years (0.3 to 5.2 years) and in 15 transgender females treated from 14.9 (± 1.9) years for 1.3 years (0.5 to 3.8 years), although not all changes were statistically significant (94). There was incomplete catch-up at age 22 years after sex hormone treatment from age 16.6 (± 1.4)

years for a median duration of 5.8 years (3.0 to 8.0 years) in transgender females and from age 16.4 (± 2.3) years for 5.4 years (2.8 to 7.8 years) in transgender males. Little is known about more prolonged use of GnRH analogs. Researchers reported normal BMD z scores at age 35 years in one individual who used GnRH analogs from age 13.7 years until age 18.6 years before initiating sex hormone treatment (65).

Additional data are available from individuals with late puberty or GnRH analog treatment of other indications. Some studies reported that men with constitutionally delayed puberty have decreased BMD in adulthood (95). However, other studies reported that these men have normal BMD (96, 97). Treating adults with GnRH analogs results in a decrease of BMD (98). In children with central precocious puberty, treatment with GnRH analogs has been found to result in a decrease of BMD during treatment by some (99) but not others (100). Studies have reported normal BMD after discontinuing therapy (69, 72, 73, 101, 102). In adolescents treated with growth hormone who are small for gestational age and have normal pubertal timing, 2-year GnRH analog treatments did not adversely affect BMD (103). Calcium supplementation may be beneficial in optimizing bone health in GnRH analog-treated individuals (104). There are no studies of vitamin D supplementation in this context, but clinicians should offer supplements to vitamin D-deficient adolescents. Physical activity, especially during growth, is important for bone mass in healthy individuals (103) and is therefore likely to be beneficial for bone health in GnRH analog-treated subjects.

GnRH analogs did not induce a change in body mass index standard deviation score in GD/gender-incongruent adolescents (94) but caused an increase in fat mass and decrease in lean body mass percentage (92). Studies in girls treated for precocious puberty also reported a stable body mass index standard deviation score during treatment (72) and body mass index and body composition comparable to controls after treatment (73).

Arterial hypertension has been reported as an adverse effect in a few girls treated with GnRH analogs for precocious/early puberty (105, 106). Blood pressure monitoring before and during treatment is recommended.

Individuals may also experience hot flashes, fatigue, and mood alterations as a consequence of pubertal suppression. There is no consensus on treatment of these side effects in this context.

It is recommended that any use of pubertal blockers (and subsequent use of sex hormones, as detailed below) include a discussion about implications for fertility (see recommendation 1.3). Transgender adolescents may

want to preserve fertility, which may be otherwise compromised if puberty is suppressed at an early stage and the individual completes phenotypic transition with the use of sex hormones.

Limited data are available regarding the effects of GnRH analogs on brain development. A single cross-sectional study demonstrated no compromise of executive function (107), but animal data suggest there may be an effect of GnRH analogs on cognitive function (108).

Values and preferences

Our recommendation of GnRH analogs places a higher value on the superior efficacy, safety, and reversibility of the pubertal hormone suppression achieved (as compared with the alternatives) and a relatively lower value on limiting the cost of therapy. Of the available alternatives, depot and oral progestin preparations are effective. Experience with this treatment dates back prior to the emergence of GnRH analogs for treating precocious puberty in papers from the 1960s and early 1970s (109–112). These compounds are usually safe, but some side effects have been reported (113–115). Only two recent studies involved transgender youth (116, 117). One of these studies described the use of oral lynestrenol monotherapy followed by the addition of testosterone treatment in transgender boys who were at Tanner stage B4 or further at the start of treatment (117). They found lynestrenol safe, but gonadotropins were not fully suppressed. The study reported metrorrhagia in approximately half of the individuals, mainly in the first 6 months. Acne, headache, hot flashes, and fatigue were other frequent side effects. Another progestin that has been studied in the United States is medroxyprogesterone. This agent is not as effective as GnRH analogs in lowering endogenous sex hormones either and may be associated with other side effects (116). Progestin preparations may be an acceptable treatment for persons without access to GnRH analogs or with a needle phobia. If GnRH analog treatment is not available (insurance denial, prohibitive cost, or other reasons), postpubertal, transgender female adolescents may be treated with an antiandrogen that directly suppresses androgen synthesis or action (see adult section).

Remarks

Measurements of gonadotropin and sex steroid levels give precise information about gonadal axis suppression, although there is insufficient evidence for any specific short-term monitoring scheme in children treated with GnRH analogs (88). If the gonadal axis is not completely suppressed—as evidenced by (for example) menses, erections, or progressive hair growth—the interval of GnRH analog treatment can be shortened or the dose increased. During treatment, adolescents should be monitored for negative effects of delaying puberty, including a halted growth spurt and impaired bone mineral accretion. Table 7 illustrates a suggested clinical protocol.

Anthropometric measurements and X-rays of the left hand to monitor bone age are informative for evaluating growth. To assess BMD, clinicians can perform dual-energy X-ray absorptiometry scans.

- 2.4. In adolescents who request sex hormone treatment (given this is a partly irreversible treatment), we recommend initiating treatment using a gradually increasing dose schedule (see Table 8) after a multidisciplinary team of medical and MHPs has confirmed the persistence of GD/gender incongruence and sufficient mental capacity to give informed consent, which most adolescents have by age 16 years (Table 5). (1 ⊕⊕○○)
- 2.5. We recognize that there may be compelling reasons to initiate sex hormone treatment prior to the age of 16 years in some adolescents with GD/gender incongruence, even though there are minimal published studies of gender-affirming hormone treatments administered before age 13.5 to 14 years. As with the care of adolescents ≥16 years of age, we recommend that an expert multidisciplinary team of medical and MHPs manage this treatment. (1 ⊕○○○)
- 2.6. We suggest monitoring clinical pubertal development every 3 to 6 months and laboratory parameters every 6 to 12 months during sex hormone treatment (Table 9). (2 ⊕⊕○○)

Table 7. Baseline and Follow-Up Protocol During Suppression of Puberty

Every 3–6 mo
Anthropometry: height, weight, sitting height, blood pressure, Tanner stages
Every 6–12 mo
Laboratory: LH, FSH, E2/T, 25OH vitamin D
Every 1–2 y
Bone density using DXA
Bone age on X-ray of the left hand (if clinically indicated)

Adapted from Hembree *et al.* (118).

Abbreviations: DXA, dual energy X ray absorptiometry; E2, estradiol; FSH, follicle stimulating hormone; LH, luteinizing hormone; T, testosterone;

Table 8. Protocol Induction of Puberty

Induction of female puberty with oral 17β -estradiol, increasing the dose every 6 mo:

5 $\mu\text{g}/\text{kg}/\text{d}$

10 $\mu\text{g}/\text{kg}/\text{d}$

15 $\mu\text{g}/\text{kg}/\text{d}$

20 $\mu\text{g}/\text{kg}/\text{d}$

Adult dose = 2–6 mg/d

In postpubertal transgender female adolescents, the dose of 17β -estradiol can be increased more rapidly:

1 mg/d for 6 mo

2 mg/d

Induction of female puberty with transdermal 17β -estradiol, increasing the dose every 6 mo (new patch is placed every 3.5 d):

6.25–12.5 $\mu\text{g}/24$ h (cut 25- μg patch into quarters, then halves)

25 $\mu\text{g}/24$ h

37.5 $\mu\text{g}/24$ h

Adult dose = 50–200 $\mu\text{g}/24$ h

For alternatives once at adult dose, see Table 11.

Adjust maintenance dose to mimic physiological estradiol levels (see Table 15).

Induction of male puberty with testosterone esters increasing the dose every 6 mo (IM or SC):

25 mg/m²/2 wk (or alternatively, half this dose weekly, or double the dose every 4 wk)

50 mg/m²/2 wk

75 mg/m²/2 wk

100 mg/m²/2 wk

Adult dose = 100–200 mg every 2 wk

In postpubertal transgender male adolescents the dose of testosterone esters can be increased more rapidly:

75 mg/2 wk for 6 mo

125 mg/2 wk

For alternatives once at adult dose, see Table 11.

Adjust maintenance dose to mimic physiological testosterone levels (see Table 14).

Adapted from Hembree et al. (118).

Abbreviations: IM, intramuscularly; SC, subcutaneously.

Evidence

Adolescents develop competence in decision making at their own pace. Ideally, the supervising medical professionals should individually assess this competence, although no objective tools to make such an assessment are currently available.

Many adolescents have achieved a reasonable level of competence by age 15 to 16 years (119), and in many countries 16-year-olds are legally competent with regard to medical decision making (120). However, others believe that although some capacities are generally achieved before age 16 years, other abilities (such as good risk

assessment) do not develop until well after 18 years (121). They suggest that health care procedures should be divided along a matrix of relative risk, so that younger adolescents can be allowed to decide about low-risk procedures, such as most diagnostic tests and common therapies, but not about high-risk procedures, such as most surgical procedures (121).

Currently available data from transgender adolescents support treatment with sex hormones starting at age 16 years (63, 122). However, some patients may incur potential risks by waiting until age 16 years. These include the potential risk to bone health if puberty is suppressed

Table 9. Baseline and Follow-up Protocol During Induction of Puberty

Every 3–6 mo

- Anthropometry: height, weight, sitting height, blood pressure, Tanner stages

Every 6–12 mo

- In transgender males: hemoglobin/hematocrit, lipids, testosterone, 25OH vitamin D
- In transgender females: prolactin, estradiol, 25OH vitamin D

Every 1–2 y

- BMD using DXA
- Bone age on X-ray of the left hand (if clinically indicated)

BMD should be monitored into adulthood (until the age of 25–30 y or until peak bone mass has been reached).

For recommendations on monitoring once pubertal induction has been completed, see Tables 14 and 15.

Adapted from Hembree et al. (118).

Abbreviation: DXA, dual energy X ray absorptiometry.

for 6 to 7 years before initiating sex hormones (*e.g.*, if someone reached Tanner stage 2 at age 9-10 years old). Additionally, there may be concerns about inappropriate height and potential harm to mental health (emotional and social isolation) if initiation of secondary sex characteristics must wait until the person has reached 16 years of age. However, only minimal data supporting earlier use of gender-affirming hormones in transgender adolescents currently exist (63). Clearly, long-term studies are needed to determine the optimal age of sex hormone treatment in GD/gender-incongruent adolescents.

The MHP who has followed the adolescent during GnRH analog treatment plays an essential role in assessing whether the adolescent is eligible to start sex hormone therapy and capable of consenting to this treatment (Table 5). Support of the family/environment is essential. Prior to the start of sex hormones, clinicians should discuss the implications for fertility (see recommendation 1.5). Throughout pubertal induction, an MHP and a pediatric endocrinologist (or other clinician competent in the evaluation and induction of pubertal development) should monitor the adolescent. In addition to monitoring therapy, it is also important to pay attention to general adolescent health issues, including healthy life style choices, such as not smoking, contraception, and appropriate vaccinations (*e.g.*, human papillomavirus).

For the induction of puberty, clinicians can use a similar dose scheme for hypogonadal adolescents with GD/gender incongruence as they use in other individuals with hypogonadism, carefully monitoring for desired and undesired effects (Table 8). In transgender female adolescents, transdermal 17β -estradiol may be an alternative for oral 17β -estradiol. It is increasingly used for pubertal induction in hypogonadal females. However, the absence of low-dose estrogen patches may be a problem. As a result, individuals may need to cut patches to size themselves to achieve appropriate dosing (123). In transgender male adolescents, clinicians can give testosterone injections intramuscularly or subcutaneously (124, 125).

When puberty is initiated with a gradually increasing schedule of sex steroid doses, the initial levels will not be high enough to suppress endogenous sex steroid secretion. Gonadotropin secretion and endogenous production of testosterone may resume and interfere with the effectiveness of estrogen treatment, in transgender female adolescents (126, 127). Therefore, continuation of GnRH analog treatment is advised until gonadectomy. Given that GD/gender-incongruent adolescents may opt not to have gonadectomy, long-term studies are necessary to examine the potential risks of prolonged GnRH analog treatment. Alternatively, in transgender male adolescents, GnRH analog treatment can be discontinued once an

adult dose of testosterone has been reached and the individual is well virilized. If uterine bleeding occurs, a progestin can be added. However, the combined use of a GnRH analog (for ovarian suppression) and testosterone may enable phenotypic transition with a lower dose of testosterone in comparison with testosterone alone. If there is a wish or need to discontinue GnRH analog treatment in transgender female adolescents, they may be treated with an antiandrogen that directly suppresses androgen synthesis or action (see section 3.0 “Hormonal Therapy for Transgender Adults”).

Values and preferences

The recommendation to initiate pubertal induction only when the individual has sufficient mental capacity (roughly age 16 years) to give informed consent for this partly irreversible treatment places a higher value on the ability of the adolescent to fully understand and oversee the partially irreversible consequences of sex hormone treatment and to give informed consent. It places a lower value on the possible negative effects of delayed puberty. We may not currently have the means to weigh adequately the potential benefits of waiting until around age 16 years to initiate sex hormones vs the potential risks/harm to BMD and the sense of social isolation from having the timing of puberty be so out of sync with peers (128).

Remarks

Before starting sex hormone treatment, effects on fertility and options for fertility preservation should be discussed. Adult height may be a concern in transgender adolescents. In a transgender female adolescent, clinicians may consider higher doses of estrogen or a more rapid tempo of dose escalation during pubertal induction. There are no established treatments yet to augment adult height in a transgender male adolescent with open epiphyses during pubertal induction. It is not uncommon for transgender adolescents to present for clinical services after having completed or nearly completed puberty. In such cases, induction of puberty with sex hormones can be done more rapidly (see Table 8). Additionally, an adult dose of testosterone in transgender male adolescents may suffice to suppress the gonadal axis without the need to use a separate agent. At the appropriate time, the multidisciplinary team should adequately prepare the adolescent for transition to adult care.

3.0 Hormonal Therapy for Transgender Adults

The two major goals of hormonal therapy are (1) to reduce endogenous sex hormone levels, and thus reduce

the secondary sex characteristics of the individual's designated gender, and (2) to replace endogenous sex hormone levels consistent with the individual's gender identity by using the principles of hormone replacement treatment of hypogonadal patients. The timing of these two goals and the age at which to begin treatment with the sex hormones of the chosen gender is codetermined in collaboration with both the person pursuing transition and the health care providers. The treatment team should include a medical provider knowledgeable in transgender hormone therapy, an MHP knowledgeable in GD/gender incongruence and the mental health concerns of transition, and a primary care provider able to provide care appropriate for transgender individuals. The physical changes induced by this sex hormone transition are usually accompanied by an improvement in mental well-being (129, 130).

- 3.1. We recommend that clinicians confirm the diagnostic criteria of GD/gender incongruence and the criteria for the endocrine phase of gender transition before beginning treatment. (1 ⊕⊕⊕⊕○)
- 3.2. We recommend that clinicians evaluate and address medical conditions that can be exacerbated by hormone depletion and treatment with sex hormones of the affirmed gender before beginning treatment (Table 10). (1 ⊕⊕⊕⊕○)
- 3.3. We suggest that clinicians measure hormone levels during treatment to ensure that endogenous sex steroids are suppressed and administered sex steroids are maintained in the normal physiologic range for the affirmed gender. (2 ⊕⊕○⊕○)

Evidence

It is the responsibility of the treating clinician to confirm that the person fulfills criteria for treatment. The treating clinician should become familiar with the terms and criteria presented in Tables 1–5 and take a thorough history from the patient in collaboration with the other members of the treatment team. The treating clinician must ensure that the desire for transition is appropriate; the consequences, risks, and benefits of treatment are well understood; and the desire for transition persists. They also need to discuss fertility preservation options (see recommendation 1.3) (67, 68).

Transgender males

Clinical studies have demonstrated the efficacy of several different androgen preparations to induce masculinization in transgender males (Appendix A) (113, 114, 131–134). Regimens to change secondary sex characteristics follow the general principle of hormone replacement treatment of male hypogonadism (135). Clinicians can use either parenteral or transdermal preparations to achieve testosterone values in the normal male range (this is dependent on the specific assay, but is typically 320 to 1000 ng/dL) (Table 11) (136). Sustained supraphysiologic levels of testosterone increase the risk of adverse reactions (see section 4.0 “Adverse Outcome Prevention and Long-Term Care”) and should be avoided.

Similar to androgen therapy in hypogonadal men, testosterone treatment in transgender males results in increased muscle mass and decreased fat mass, increased facial hair and acne, male pattern baldness in those genetically predisposed, and increased sexual desire (137).

Table 10. Medical Risks Associated With Sex Hormone Therapy

Transgender female: estrogen

Very high risk of adverse outcomes:

- Thromboembolic disease

Moderate risk of adverse outcomes:

- Macroprolactinoma
- Breast cancer
- Coronary artery disease
- Cerebrovascular disease
- Cholelithiasis
- Hypertriglyceridemia

Transgender male: testosterone

Very high risk of adverse outcomes:

- Erythrocytosis (hematocrit > 50%)

Moderate risk of adverse outcomes:

- Severe liver dysfunction (transaminases > threefold upper limit of normal)
- Coronary artery disease
- Cerebrovascular disease
- Hypertension
- Breast or uterine cancer

Table 11. Hormone Regimens in Transgender Persons

Transgender females ^a	
Estrogen	
Oral	
Estradiol	2.0–6.0 mg/d
Transdermal	
Estradiol transdermal patch (New patch placed every 3–5 d)	0.025–0.2 mg/d
Parenteral	
Estradiol valerate or cypionate	5–30 mg IM every 2 wk 2–10 mg IM every week
Anti-androgens	
Spironolactone	100–300 mg/d
Cyproterone acetate ^b	25–50 mg/d
GnRH agonist	3.75 mg SQ (SC) monthly 11.25 mg SQ (SC) 3-monthly
Transgender males	
Testosterone	
Parenteral testosterone	
Testosterone enanthate or cypionate	100–200 mg SQ (IM) every 2 wk or SQ (SC) 50% per week
Testosterone undecanoate ^c	1000 mg every 12 wk
Transdermal testosterone	
Testosterone gel 1.6% ^d	50–100 mg/d
Testosterone transdermal patch	2.5–7.5 mg/d

Abbreviations: IM, intramuscularly; SQ, sequentially; SC, subcutaneously.

^aEstrogens used with or without antiandrogens or GnRH agonist.

^bNot available in the United States.

^cOne thousand milligrams initially followed by an injection at 6 wk then at 12 wk intervals.

^dAvoid cutaneous transfer to other individuals.

In transgender males, testosterone will result in clitoromegaly, temporary or permanent decreased fertility, deepening of the voice, cessation of menses (usually), and a significant increase in body hair, particularly on the face, chest, and abdomen. Cessation of menses may occur within a few months with testosterone treatment alone, although high doses of testosterone may be required. If uterine bleeding continues, clinicians may consider the addition of a progestational agent or endometrial ablation (138). Clinicians may also administer GnRH analogs or depot medroxyprogesterone to stop menses prior to testosterone treatment.

Transgender females

The hormone regimen for transgender females is more complex than the transgender male regimen (Appendix B). Treatment with physiologic doses of estrogen alone is insufficient to suppress testosterone levels into the normal range for females (139). Most published clinical studies report the need for adjunctive therapy to achieve testosterone levels in the female range (21, 113, 114, 132–134, 139, 140).

Multiple adjunctive medications are available, such as progestins with antiandrogen activity and GnRH agonists (141). Spironolactone works by directly blocking androgens during their interaction with the androgen

receptor (114, 133, 142). It may also have estrogenic activity (143). Cyproterone acetate, a progestational compound with antiandrogenic properties (113, 132, 144), is widely used in Europe. 5 α -Reductase inhibitors do not reduce testosterone levels and have adverse effects (145).

Dittrich *et al.* (141) reported that monthly doses of the GnRH agonist goserelin acetate in combination with estrogen were effective in reducing testosterone levels with a low incidence of adverse reactions in 60 transgender females. Leuprolide and transdermal estrogen were as effective as cyproterone and transdermal estrogen in a comparative retrospective study (146).

Patients can take estrogen as oral conjugated estrogens, oral 17 β -estradiol, or transdermal 17 β -estradiol. Among estrogen options, the increased risk of thromboembolic events associated with estrogens in general seems most concerning with ethinyl estradiol specifically (134, 140, 141), which is why we specifically suggest that it not be used in any transgender treatment plan. Data distinguishing among other estrogen options are less well established although there is some thought that oral routes of administration are more thrombogenic due to the “first pass effect” than are transdermal and parenteral routes, and that the risk of thromboembolic events is dose-dependent. Injectable estrogen and sublingual

estrogen may benefit from avoiding the first pass effect, but they can result in more rapid peaks with greater overall periodicity and thus are more difficult to monitor (147, 148). However, there are no data demonstrating that increased periodicity is harmful otherwise.

Clinicians can use serum estradiol levels to monitor oral, transdermal, and intramuscular estradiol. Blood tests cannot monitor conjugated estrogens or synthetic estrogen use. Clinicians should measure serum estradiol and serum testosterone and maintain them at the level for premenopausal females (100 to 200 pg/mL and <50 ng/dL, respectively). The transdermal preparations and injectable estradiol cypionate or valerate preparations may confer an advantage in older transgender females who may be at higher risk for thromboembolic disease (149).

Values

Our recommendation to maintain levels of gender-affirming hormones in the normal adult range places a high value on the avoidance of the long-term complications of pharmacologic doses. Those patients receiving endocrine treatment who have relative contraindications to hormones should have an in-depth discussion with their physician to balance the risks and benefits of therapy.

Remarks

Clinicians should inform all endocrine-treated individuals of all risks and benefits of gender-affirming hormones prior to initiating therapy. Clinicians should strongly encourage tobacco use cessation in transgender females to avoid increased risk of VTE and cardiovascular complications. We strongly discourage the unsupervised use of hormone therapy (150).

Not all individuals with GD/gender incongruence seek treatment as described (*e.g.*, male-to-eunuchs and individuals seeking partial transition). Tailoring current protocols to the individual may be done within the context of accepted safety guidelines using a multidisciplinary approach including mental health. No evidence-based protocols are available for these groups (151). We need prospective studies to better understand treatment options for these persons.

- 3.4. We suggest that endocrinologists provide education to transgender individuals undergoing treatment about the onset and time course of physical changes induced by sex hormone treatment. (2 ⊕○○○)

Evidence

Transgender males

Physical changes that are expected to occur during the first 1 to 6 months of testosterone therapy include

cessation of menses, increased sexual desire, increased facial and body hair, increased oiliness of skin, increased muscle, and redistribution of fat mass. Changes that occur within the first year of testosterone therapy include deepening of the voice (152, 153), clitoromegaly, and male pattern hair loss (in some cases) (114, 144, 154, 155) (Table 12).

Transgender females

Physical changes that may occur in transgender females in the first 3 to 12 months of estrogen and anti-androgen therapy include decreased sexual desire, decreased spontaneous erections, decreased facial and body hair (usually mild), decreased oiliness of skin, increased breast tissue growth, and redistribution of fat mass (114, 139, 149, 154, 155, 161) (Table 13). Breast development is generally maximal at 2 years after initiating hormones (114, 139, 149, 155). Over a long period of time, the prostate gland and testicles will undergo atrophy.

Although the time course of breast development in transgender females has been studied (150), precise information about other changes induced by sex hormones is lacking (141). There is a great deal of variability among individuals, as evidenced during pubertal development. We all know that a major concern for transgender females is breast development. If we work with estrogens, the result will be often not what the transgender female expects.

Alternatively, there are transgender females who report an anecdotal improved breast development, mood, or sexual desire with the use of progestogens. However, there have been no well-designed studies of the role of progestogens in feminizing hormone regimens, so the question is still open.

Our knowledge concerning the natural history and effects of different cross-sex hormone therapies on breast

Table 12. Masculinizing Effects in Transgender Males

Effect	Onset	Maximum
Skin oiliness/acne	1–6 mo	1–2 y
Facial/body hair growth	6–12 mo	4–5 y
Scalp hair loss	6–12 mo	— ^a
Increased muscle mass/strength	6–12 mo	2–5 y
Fat redistribution	1–6 mo	2–5 y
Cessation of menses	1–6 mo	— ^b
Clitoral enlargement	1–6 mo	1–2 y
Vaginal atrophy	1–6 mo	1–2 y
Deepening of voice	6–12 mo	1–2 y

Estimates represent clinical observations: Toorians *et al.* (149), Assche man *et al.* (156), Gooren *et al.* (157), Wierckx *et al.* (158).

^aPrevention and treatment as recommended for biological men.

^bMenorrhagia requires diagnosis and treatment by a gynecologist.

Table 13. Feminizing Effects in Transgender Females

Effect	Onset	Maximum
Redistribution of body fat	3–6 mo	2–3 y
Decrease in muscle mass and strength	3–6 mo	1–2 y
Softening of skin/decreased oiliness	3–6 mo	Unknown
Decreased sexual desire	1–3 mo	3–6 mo
Decreased spontaneous erections	1–3 mo	3–6 mo
Male sexual dysfunction	Variable	Variable
Breast growth	3–6 mo	2–3 y
Decreased testicular volume	3–6 mo	2–3 y
Decreased sperm production	Unknown	>3 y ^a
Decreased terminal hair growth	6–12 mo	>3 y ^a
Scalp hair	Variable	— ^b
Voice changes	None	— ^c

Estimates represent clinical observations: Toorians *et al.* (149), Asscheman *et al.* (156), Gooren *et al.* (157).

^aComplete removal of male sexual hair requires electrolysis or laser treatment or both.

^bFamilial scalp hair loss may occur if estrogens are stopped.

^cTreatment by speech pathologists for voice training is most effective.

development in transgender females is extremely sparse and based on the low quality of evidence. Current evidence does not indicate that progestogens enhance breast development in transgender females, nor does evidence prove the absence of such an effect. This prevents us from drawing any firm conclusion at this moment and demonstrates the need for further research to clarify these important clinical questions (162).

Values and preferences

Transgender persons have very high expectations regarding the physical changes of hormone treatment and are aware that body changes can be enhanced by surgical procedures (*e.g.*, breast, face, and body habitus). Clear expectations for the extent and timing of sex hormone-induced changes may prevent the potential harm and expense of unnecessary procedures.

4.0 Adverse Outcome Prevention and Long-Term Care

Hormone therapy for transgender males and females confers many of the same risks associated with sex hormone replacement therapy in nontransgender persons. The risks arise from and are worsened by inadvertent or intentional use of supraphysiologic doses of sex hormones, as well as use of inadequate doses of sex hormones to maintain normal physiology (131, 139).

- 4.1. We suggest regular clinical evaluation for physical changes and potential adverse changes in response to sex steroid hormones and laboratory monitoring of sex steroid hormone levels every

3 months during the first year of hormone therapy for transgender males and females and then once or twice yearly. (2 ⊕⊕○○)

Evidence

Pretreatment screening and appropriate regular medical monitoring are recommended for both transgender males and females during the endocrine transition and periodically thereafter (26, 155). Clinicians should monitor weight and blood pressure, conduct physical exams, and assess routine health questions, such as tobacco use, symptoms of depression, and risk of adverse events such as deep vein thrombosis/pulmonary embolism and other adverse effects of sex steroids.

Transgender males

Table 14 contains a standard monitoring plan for transgender males on testosterone therapy (154, 159). Key issues include maintaining testosterone levels in the physiologic normal male range and avoiding adverse events resulting from excess testosterone therapy, particularly erythrocytosis, sleep apnea, hypertension, excessive weight gain, salt retention, lipid changes, and excessive or cystic acne (135).

Because oral 17-alkylated testosterone is not recommended, serious hepatic toxicity is not anticipated with parenteral or transdermal testosterone use (163, 164). Past concerns regarding liver toxicity with testosterone have been alleviated with subsequent reports that indicate the risk of serious liver disease is minimal (144, 165, 166).

Transgender females

Table 15 contains a standard monitoring plan for transgender females on estrogens, gonadotropin suppression, or antiandrogens (160). Key issues include avoiding supraphysiologic doses or blood levels of estrogen that may lead to increased risk for thromboembolic disease, liver dysfunction, and hypertension. Clinicians should monitor serum estradiol levels using laboratories participating in external quality control, as measurements of estradiol in blood can be very challenging (167).

VTE may be a serious complication. A study reported a 20-fold increase in venous thromboembolic disease in a large cohort of Dutch transgender subjects (161). This increase may have been associated with the use of the synthetic estrogen, ethinyl estradiol (149). The incidence decreased when clinicians stopped administering ethinyl estradiol (161). Thus, the use of synthetic estrogens and conjugated estrogens is undesirable because of the inability to regulate doses by measuring serum levels and the risk of thromboembolic disease. In a German gender clinic, deep vein thrombosis occurred in 1 of 60 of transgender females treated with a GnRH analog and oral

Table 14. Monitoring of Transgender Persons on Gender-Affirming Hormone Therapy: Transgender Male

1. Evaluate patient every 3 mo in the first year and then one to two times per year to monitor for appropriate signs of virilization and for development of adverse reactions.
2. Measure serum testosterone every 3 mo until levels are in the normal physiologic male range:^a
 - a. For testosterone enanthate/cypionate injections, the testosterone level should be measured midway between injections. The target level is 400–700 ng/dL to 400 ng/dL. Alternatively, measure peak and trough levels to ensure levels remain in the normal male range.
 - b. For parenteral testosterone undecanoate, testosterone should be measured just before the following injection. If the level is <400 ng/dL, adjust dosing interval.
 - c. For transdermal testosterone, the testosterone level can be measured no sooner than after 1 wk of daily application (at least 2 h after application).
3. Measure hematocrit or hemoglobin at baseline and every 3 mo for the first year and then one to two times a year. Monitor weight, blood pressure, and lipids at regular intervals.
4. Screening for osteoporosis should be conducted in those who stop testosterone treatment, are not compliant with hormone therapy, or who develop risks for bone loss.
5. If cervical tissue is present, monitoring as recommended by the American College of Obstetricians and Gynecologists.
6. Ovariectomy can be considered after completion of hormone transition.
7. Conduct sub- and periareolar annual breast examinations if mastectomy performed. If mastectomy is not performed, then consider mammograms as recommended by the American Cancer Society.

^aAdapted from Lapauw *et al.* (154) and Ott *et al.* (159).

estradiol (141). The patient who developed a deep vein thrombosis was found to have a homozygous C677 T mutation in the methylenetetrahydrofolate reductase gene. In an Austrian gender clinic, administering gender-affirming hormones to 162 transgender females and 89 transgender males was not associated with VTE, despite an 8.0% and 5.6% incidence of thrombophilia (159). A more recent multinational study reported only 10 cases of VTE from a cohort of 1073 subjects (168). Thrombophilia screening of transgender persons initiating hormone treatment should be restricted to those with a personal or family history of VTE (159). Monitoring D-dimer levels during treatment is not recommended (169).

- 4.2. We suggest periodically monitoring prolactin levels in transgender females treated with estrogens. (2 ⊕⊕○○)

Evidence

Estrogen therapy can increase the growth of pituitary lactotroph cells. There have been several reports of prolactinomas occurring after long-term, high-dose

estrogen therapy (170–173). Up to 20% of transgender females treated with estrogens may have elevations in prolactin levels associated with enlargement of the pituitary gland (156). In most cases, the serum prolactin levels will return to the normal range with a reduction or discontinuation of the estrogen therapy or discontinuation of cyproterone acetate (157, 174, 175).

The onset and time course of hyperprolactinemia during estrogen treatment are not known. Clinicians should measure prolactin levels at baseline and then at least annually during the transition period and every 2 years thereafter. Given that only a few case studies reported prolactinomas, and prolactinomas were not reported in large cohorts of estrogen-treated persons, the risk is likely to be very low. Because the major presenting findings of microprolactinomas (hypogonadism and sometimes gynecomastia) are not apparent in transgender females, clinicians may perform radiologic examinations of the pituitary in those patients whose prolactin levels persistently increase despite stable or reduced estrogen levels. Some transgender individuals receive psychotropic medications that can increase prolactin levels (174).

Table 15. Monitoring of Transgender Persons on Gender-Affirming Hormone Therapy: Transgender Female

1. Evaluate patient every 3 mo in the first year and then one to two times per year to monitor for appropriate signs of feminization and for development of adverse reactions.
2. Measure serum testosterone and estradiol every 3 mo.
 - a. Serum testosterone levels should be <50 ng/dL.
 - b. Serum estradiol should not exceed the peak physiologic range: 100–200 pg/mL.
3. For individuals on spironolactone, serum electrolytes, particularly potassium, should be monitored every 3 mo in the first year and annually thereafter.
4. Routine cancer screening is recommended, as in nontransgender individuals (all tissues present).
5. Consider BMD testing at baseline (160). In individuals at low risk, screening for osteoporosis should be conducted at age 60 years or in those who are not compliant with hormone therapy.

This table presents strong recommendations and does not include lower level recommendations.

- 4.3. We suggest that clinicians evaluate transgender persons treated with hormones for cardiovascular risk factors using fasting lipid profiles, diabetes screening, and/or other diagnostic tools. (2 ⊕⊕○○)

Evidence

Transgender males

Administering testosterone to transgender males results in a more atherogenic lipid profile with lowered high-density lipoprotein cholesterol and higher triglyceride and low-density lipoprotein cholesterol values (176–179). Studies of the effect of testosterone on insulin sensitivity have mixed results (178, 180). A randomized, open-label uncontrolled safety study of transgender males treated with testosterone undecanoate demonstrated no insulin resistance after 1 year (181, 182). Numerous studies have demonstrated the effects of sex hormone treatment on the cardiovascular system (160, 179, 183, 184). Long-term studies from The Netherlands found no increased risk for cardiovascular mortality (161). Likewise, a meta-analysis of 19 randomized trials in nontransgender males on testosterone replacement showed no increased incidence of cardiovascular events (185). A systematic review of the literature found that data were insufficient (due to very low-quality evidence) to allow a meaningful assessment of patient-important outcomes, such as death, stroke, myocardial infarction, or VTE in transgender males (176). Future research is needed to ascertain the potential harm of hormonal therapies (176). Clinicians should manage cardiovascular risk factors as they emerge according to established guidelines (186).

Transgender females

A prospective study of transgender females found favorable changes in lipid parameters with increased high-density lipoprotein and decreased low-density lipoprotein concentrations (178). However, increased weight, blood pressure, and markers of insulin resistance attenuated these favorable lipid changes. In a meta-analysis, only serum triglycerides were higher at ≥24 months without changes in other parameters (187). The largest cohort of transgender females (mean age 41 years, followed for a mean of 10 years) showed no increase in cardiovascular mortality despite a 32% rate of tobacco use (161).

Thus, there is limited evidence to determine whether estrogen is protective or detrimental on lipid and glucose metabolism in transgender females (176). With aging, there is usually an increase of body weight. Therefore, as with nontransgender individuals, clinicians should

monitor and manage glucose and lipid metabolism and blood pressure regularly according to established guidelines (186).

- 4.4. We recommend that clinicians obtain BMD measurements when risk factors for osteoporosis exist, specifically in those who stop sex hormone therapy after gonadectomy. (1 ⊕⊕○○)

Evidence

Transgender males

Baseline bone mineral measurements in transgender males are generally in the expected range for their pre-treatment gender (188). However, adequate dosing of testosterone is important to maintain bone mass in transgender males (189, 190). In one study (190), serum LH levels were inversely related to BMD, suggesting that low levels of sex hormones were associated with bone loss. Thus, LH levels in the normal range may serve as an indicator of the adequacy of sex steroid administration to preserve bone mass. The protective effect of testosterone may be mediated by peripheral conversion to estradiol, both systemically and locally in the bone.

Transgender females

A baseline study of BMD reported T scores less than -2.5 in 16% of transgender females (191). In aging males, studies suggest that serum estradiol more positively correlates with BMD than does testosterone (192, 193) and is more important for peak bone mass (194). Estrogen preserves BMD in transgender females who continue on estrogen and antiandrogen therapies (188, 190, 191, 195, 196).

Fracture data in transgender males and females are not available. Transgender persons who have undergone gonadectomy may choose not to continue consistent sex steroid treatment after hormonal and surgical sex reassignment, thereby becoming at risk for bone loss. There have been no studies to determine whether clinicians should use the sex assigned at birth or affirmed gender for assessing osteoporosis (e.g., when using the FRAX tool). Although some researchers use the sex assigned at birth (with the assumption that bone mass has usually peaked for transgender people who initiate hormones in early adulthood), this should be assessed on a case-by-case basis until there are more data available. This assumption will be further complicated by the increasing prevalence of transgender people who undergo hormonal transition at a pubertal age or soon after puberty. Sex for comparison within risk assessment tools may be based on the age at which hormones were initiated and the length of exposure to hormones. In some cases, it may be

reasonable to assess risk using both the male and female calculators and using an intermediate value. Because all subjects underwent normal pubertal development, with known effects on bone size, reference values for birth sex were used for all participants (154).

- 4.5. We suggest that transgender females with no known increased risk of breast cancer follow breast-screening guidelines recommended for those designated female at birth. (2 ⊕⊕○○)
- 4.6. We suggest that transgender females treated with estrogens follow individualized screening according to personal risk for prostatic disease and prostate cancer. (2 ⊕○○○)

Evidence

Studies have reported a few cases of breast cancer in transgender females (197–200). A Dutch study of 1800 transgender females followed for a mean of 15 years (range of 1–30 years) found one case of breast cancer. The Women's Health Initiative study reported that females taking conjugated equine estrogen without progesterone for 7 years did not have an increased risk of breast cancer as compared with females taking placebo (137).

In transgender males, a large retrospective study conducted at the U.S. Veterans Affairs medical health system identified seven breast cancers (194). The authors reported that this was not above the expected rate of breast cancers in cisgender females in this cohort. Furthermore, they did report one breast cancer that developed in a transgender male patient after mastectomy, supporting the fact that breast cancer can occur even after mastectomy. Indeed, there have been case reports of breast cancer developing in subareolar tissue in transgender males, which occurred after mastectomy (201, 202).

Women with primary hypogonadism (Turner syndrome) treated with estrogen replacement exhibited a significantly decreased incidence of breast cancer as compared with national standardized incidence ratios (203, 204). These studies suggest that estrogen therapy does not increase the risk of breast cancer in the short term (<20 to 30 years). We need long-term studies to determine the actual risk, as well as the role of screening mammograms. Regular examinations and gynecologic advice should determine monitoring for breast cancer.

Prostate cancer is very rare before the age of 40, especially with androgen deprivation therapy (205). Childhood or pubertal castration results in regression of the prostate and adult castration reverses benign prostate hypertrophy (206). Although van Kesteren *et al.* (207) reported that estrogen therapy does not induce hypertrophy or premalignant changes in the prostates of

transgender females, studies have reported cases of benign prostatic hyperplasia in transgender females treated with estrogens for 20 to 25 years (208, 209). Studies have also reported a few cases of prostate carcinoma in transgender females (210–214).

Transgender females may feel uncomfortable scheduling regular prostate examinations. Gynecologists are not trained to screen for prostate cancer or to monitor prostate growth. Thus, it may be reasonable for transgender females who transitioned after age 20 years to have annual screening digital rectal examinations after age 50 years and prostate-specific antigen tests consistent with U.S. Preventive Services Task Force Guidelines (215).

- 4.7. We advise that clinicians determine the medical necessity of including a total hysterectomy and oophorectomy as part of gender-affirming surgery. (Ungraded Good Practice Statement)

Evidence

Although aromatization of testosterone to estradiol in transgender males has been suggested as a risk factor for endometrial cancer (216), no cases have been reported. When transgender males undergo hysterectomy, the uterus is small and there is endometrial atrophy (217, 218). Studies have reported cases of ovarian cancer (219, 220). Although there is limited evidence for increased risk of reproductive tract cancers in transgender males, health care providers should determine the medical necessity of a laparoscopic total hysterectomy as part of a gender-affirming surgery to prevent reproductive tract cancer (221).

Values

Given the discomfort that transgender males experience accessing gynecologic care, our recommendation for the medical necessity of total hysterectomy and oophorectomy places a high value on eliminating the risks of female reproductive tract disease and cancer and a lower value on avoiding the risks of these surgical procedures (related to the surgery and to the potential undesirable health consequences of oophorectomy) and their associated costs.

Remarks

The sexual orientation and type of sexual practices will determine the need and types of gynecologic care required following transition. Additionally, in certain countries, the approval required to change the sex in a birth certificate for transgender males may be dependent on having a complete hysterectomy. Clinicians should help patients research nonmedical administrative criteria and

provide counseling. If individuals decide not to undergo hysterectomy, screening for cervical cancer is the same as all other females.

5.0 Surgery for Sex Reassignment and Gender Confirmation

For many transgender adults, genital gender-affirming surgery may be the necessary step toward achieving their ultimate goal of living successfully in their desired gender role. The type of surgery falls into two main categories: (1) those that directly affect fertility and (2) those that do not. Those that change fertility (previously called sex reassignment surgery) include genital surgery to remove the penis and gonads in the male and removal of the uterus and gonads in the female. The surgeries that effect fertility are often governed by the legal system of the state or country in which they are performed. Other gender-conforming surgeries that do not directly affect fertility are not so tightly governed.

Gender-affirming surgical techniques have improved markedly during the past 10 years. Reconstructive genital surgery that preserves neurologic sensation is now the standard. The satisfaction rate with surgical reassignment of sex is now very high (187). Additionally, the mental health of the individual seems to be improved by participating in a treatment program that defines a pathway of gender-affirming treatment that includes hormones and surgery (130, 144) (Table 16).

Surgery that affects fertility is irreversible. The World Professional Association for Transgender Health Standards of Care (222) emphasizes that the “threshold of 18 should not be seen as an indication in itself for active intervention.” If the social transition has not been satisfactory, if the person is not satisfied with or is ambivalent about the effects of sex hormone treatment, or if the person is ambivalent about surgery then the individual should not be referred for surgery (223, 224).

Gender-affirming genital surgeries for transgender females that affect fertility include gonadectomy, penectomy, and creation of a neovagina (225, 226). Surgeons often invert the skin of the penis to form the wall of the vagina, and several literatures reviews have

reported on outcomes (227). Sometimes there is inadequate tissue to form a full neovagina, so clinicians have revisited using intestine and found it to be successful (87, 228, 229). Some newer vaginoplasty techniques may involve autologous oral epithelial cells (230, 231).

The scrotum becomes the labia majora. Surgeons use reconstructive surgery to fashion the clitoris and its hood, preserving the neurovascular bundle at the tip of the penis as the neurosensory supply to the clitoris. Some surgeons are also creating a sensate pedicled-spot adding a G spot to the neovagina to increase sensation (232). Most recently, plastic surgeons have developed techniques to fashion labia minora. To further complete the feminization, uterine transplants have been proposed and even attempted (233).

Neovaginal prolapse, rectovaginal fistula, delayed healing, vaginal stenosis, and other complications do sometimes occur (234, 235). Clinicians should strongly remind the transgender person to use their dilators to maintain the depth and width of the vagina throughout the postoperative period. Genital sexual responsiveness and other aspects of sexual function are usually preserved following genital gender-affirming surgery (236, 237).

Ancillary surgeries for more feminine or masculine appearance are not within the scope of this guideline. Voice therapy by a speech language pathologist is available to transform speech patterns to the affirmed gender (148). Spontaneous voice deepening occurs during testosterone treatment of transgender males (152, 238). No studies have compared the effectiveness of speech therapy, laryngeal surgery, or combined treatment.

Breast surgery is a good example of gender-confirming surgery that does not affect fertility. In all females, breast size exhibits a very broad spectrum. For transgender females to make the best informed decision, clinicians should delay breast augmentation surgery until the patient has completed at least 2 years of estrogen therapy, because the breasts continue to grow during that time (141, 155).

Another major procedure is the removal of facial and masculine-appearing body hair using either electrolysis or

Table 16. Criteria for Gender-Affirming Surgery, Which Affects Fertility

1. Persistent, well-documented gender dysphoria
2. Legal age of majority in the given country
3. Having continuously and responsibly used gender-affirming hormones for 12 mo (if there is no medical contraindication to receiving such therapy)
4. Successful continuous full-time living in the new gender role for 12 mo
5. If significant medical or mental health concerns are present, they must be well controlled
6. Demonstrable knowledge of all practical aspects of surgery (e.g., cost, required lengths of hospitalizations, likely complications, postsurgical rehabilitation)

laser treatments. Other feminizing surgeries, such as that to feminize the face, are now becoming more popular (239–241).

In transgender males, clinicians usually delay gender-affirming genital surgeries until after a few years of androgen therapy. Those surgeries that affect fertility in this group include oophorectomy, vaginectomy, and complete hysterectomy. Surgeons can safely perform them vaginally with laparoscopy. These are sometimes done in conjunction with the creation of a neopenis. The cosmetic appearance of a neopenis is now very good, but the surgery is multistage and very expensive (242, 243). Radial forearm flap seems to be the most satisfactory procedure (228, 244). Other flaps also exist (245). Surgeons can make neopenile erections possible by reinnervation of the flap and subsequent contraction of the muscle, leading to stiffening of the neopenis (246, 247), but results are inconsistent (248). Surgeons can also stiffen the penis by imbedding some mechanical device (*e.g.*, a rod or some inflatable apparatus) (249, 250). Because of these limitations, the creation of a neopenis has often been less than satisfactory. Recently, penis transplants are being proposed (233).

In fact, most transgender males do not have any external genital surgery because of the lack of access, high cost, and significant potential complications. Some choose a metaoidioplasty that brings forward the clitoris, thereby allowing them to void in a standing position without wetting themselves (251, 252). Surgeons can create the scrotum from the labia majora with good cosmetic effect and can implant testicular prostheses (253).

The most important masculinizing surgery for the transgender male is mastectomy, and it does not affect fertility. Breast size only partially regresses with androgen therapy (155). In adults, discussions about mastectomy usually take place after androgen therapy has started. Because some transgender male adolescents present after significant breast development has occurred, they may also consider mastectomy 2 years after they begin androgen therapy and before age 18 years. Clinicians should individualize treatment based on the physical and mental health status of the individual. There are now newer approaches to mastectomy with better outcomes (254, 255). These often involve chest contouring (256). Mastectomy is often necessary for living comfortably in the new gender (256).

- 5.1. We recommend that a patient pursue genital gender-affirming surgery only after the MHP and the clinician responsible for endocrine transition therapy both agree that surgery is medically

necessary and would benefit the patient's overall health and/or well-being. (1 ⊕⊕○○)

- 5.2. We advise that clinicians approve genital gender-affirming surgery only after completion of at least 1 year of consistent and compliant hormone treatment, unless hormone therapy is not desired or medically contraindicated. (Ungraded Good Practice Statement)
- 5.3. We advise that the clinician responsible for endocrine treatment and the primary care provider ensure appropriate medical clearance of transgender individuals for genital gender-affirming surgery and collaborate with the surgeon regarding hormone use during and after surgery. (Ungraded Good Practice Statement)
- 5.4. We recommend that clinicians refer hormone-treated transgender individuals for genital surgery when: (1) the individual has had a satisfactory social role change, (2) the individual is satisfied about the hormonal effects, and (3) the individual desires definitive surgical changes. (1 ⊕○○○)
- 5.5. We suggest that clinicians delay gender-affirming genital surgery involving gonadectomy and/or hysterectomy until the patient is at least 18 years old or legal age of majority in his or her country. (2 ⊕⊕○○)
- 5.6. We suggest that clinicians determine the timing of breast surgery for transgender males based upon the physical and mental health status of the individual. There is insufficient evidence to recommend a specific age requirement. (2 ⊕○○○)

Evidence

Owing to the lack of controlled studies, incomplete follow-up, and lack of valid assessment measures, evaluating various surgical approaches and techniques is difficult. However, one systematic review including a large numbers of studies reported satisfactory cosmetic and functional results for vaginoplasty/neovagina construction (257). For transgender males, the outcomes are less certain. However, the problems are now better understood (258). Several postoperative studies report significant long-term psychological and psychiatric pathology (259–261). One study showed satisfaction with breasts, genitals, and femininity increased significantly and showed the importance of surgical treatment as a key therapeutic option for transgender females (262). Another analysis demonstrated that, despite the young average age at death following surgery and the relatively larger number of individuals with somatic morbidity, the study does not allow for determination of

causal relationships between, for example, specific types of hormonal or surgical treatment received and somatic morbidity and mortality (263). Reversal surgery in regretful male-to-female transsexuals after sexual reassignment surgery represents a complex, multistage procedure with satisfactory outcomes. Further insight into the characteristics of persons who regret their decision postoperatively would facilitate better future selection of applicants eligible for sexual reassignment surgery. We need more studies with appropriate controls that examine long-term quality of life, psychosocial outcomes, and psychiatric outcomes to determine the long-term benefits of surgical treatment.

When a transgender individual decides to have gender-affirming surgery, both the hormone prescribing clinician and the MHP must certify that the patient satisfies criteria for gender-affirming surgery (Table 16).

There is some concern that estrogen therapy may cause an increased risk for venous thrombosis during or following surgery (176). For this reason, the surgeon and the hormone-prescribing clinician should collaborate in making a decision about the use of hormones before and following surgery. One study suggests that preoperative factors (such as compliance) are less important for patient satisfaction than are the physical postoperative results (56). However, other studies and clinical experience dictate that individuals who do not follow medical instructions and do not work with their physicians toward a common goal do not achieve treatment goals (264) and experience higher rates of postoperative infections and other complications (265, 266). It is also important that the person requesting surgery feels comfortable with the anatomical changes that have occurred during hormone therapy. Dissatisfaction with social and physical outcomes during the hormone transition may be a contraindication to surgery (223).

An endocrinologist or experienced medical provider should monitor transgender individuals after surgery. Those who undergo gonadectomy will require hormone replacement therapy, surveillance, or both to prevent adverse effects of chronic hormone deficiency.

Financial Disclosures of the Task Force*

Wylie C. Hembree (chair)—financial or business/organizational interests: none declared, significant financial interest or leadership position: none declared. **Peggy T. Cohen-Kettenis**—financial or business/organizational interests: none declared, significant financial interest or leadership position: none declared. **Louis Gooren**—financial or business/organizational interests: none declared, significant financial

interest or leadership position: none declared. **Sabine E. Hannema**—financial or business/organizational interests: none declared, significant financial interest or leadership position: Ferring Pharmaceuticals Inc. (lecture/conference), Pfizer (lecture). **Walter J. Meyer**—financial or business/organizational interests: none declared, significant financial interest or leadership position: none declared. **M. Hassan Murad****—financial or business/organizational interests: Mayo Clinic, Evidence-based Practice Center, significant financial interest or leadership position: none declared. **Stephen M. Rosenthal**—financial or business/organizational interests: AbbVie (consultant), National Institutes of Health (grantee), significant financial interest or leadership position: Pediatric Endocrine Society (immediate past president). **Joshua D. Safer, FACP**—financial or business/organizational interests: none declared, significant financial interest or leadership position: none declared. **Vin Tangpricha**—financial or business/organizational interests: Cystic Fibrosis Foundation (grantee), National Institutes of Health (grantee), significant financial interest or leadership position, Elsevier *Journal of Clinical and Translational Endocrinology* (editor). **Guy G. T'Sjoen**—financial or business/organizational interests: none declared, significant financial interest or leadership position: none declared.* Financial, business, and organizational disclosures of the task force cover the year prior to publication. Disclosures prior to this time period are archived.**Evidence-based reviews for this guideline were prepared under contract with the Endocrine Society.

Acknowledgments

Correspondence and Reprint Requests: The Endocrine Society, 2055 L Street NW, Suite 600, Washington, DC 20036. E mail: publications@endocrine.org; Phone: 202971 3636.

Disclosure Summary: See Financial Disclosures.

Disclaimer: The Endocrine Society's clinical practice guidelines are developed to be of assistance to endocrinologists by providing guidance and recommendations for particular areas of practice. The guidelines should not be considered inclusive of all proper approaches or methods, or exclusive of others. The guidelines cannot guarantee any specific outcome, nor do they establish a standard of care. The guidelines are not intended to dictate the treatment of a particular patient. Treatment decisions must be made based on the independent judgement of healthcare providers and each patient's individual circumstances.

The Endocrine Society makes no warranty, express or implied, regarding the guidelines and specifically excludes any warranties of merchantability and fitness for a particular use or purpose. The Society shall not be liable for direct, indirect,

special, incidental, or consequential damages related to the use of the information contained herein.

References

- Atkins D, Best D, Briss PA, Eccles M, Falck Ytter Y, Flottorp S, Guyatt GH, Harbour RT, Haugh MC, Henry D, Hill S, Jaeschke R, Leng G, Liberati A, Magrini N, Mason J, Middleton P, Mrukowicz J, O'Connell D, Oxman AD, Phillips B, Schunemann HJ, Edejer T, Varonen H, Vist GE, Williams JW, Jr, Zaza S; GRADE Working Group. Grading quality of evidence and strength of recommendations. *BMJ*. 2004;328(7454):1490.
- Swiglo BA, Murad MH, Schunemann HJ, Kunz R, Vigersky RA, Guyatt GH, Montori VM. A case for clarity, consistency, and helpfulness: state of the art clinical practice guidelines in endocrinology using the grading of recommendations, assessment, development, and evaluation system. *J Clin Endocrinol Metab*. 2008;93(3):666–673.
- Bullough VL. Transsexualism in history. *Arch Sex Behav*. 1975;4(5):561–571.
- Benjamin H. The transsexual phenomenon. *Trans N Y Acad Sci*. 1967;29(4):428–430.
- Meyerowitz J. *How Sex Changed: A History of Transsexuality in the United States*. Cambridge, MA: Harvard University Press; 2002.
- Hirschfeld M. *Was muss das Volk vom Dritten Geschlecht wissen*. Verlag Max Spohr, Leipzig; 1901.
- Fisk NM. Editorial: Gender dysphoria syndrome—the conceptualization that liberalizes indications for total gender reorientation and implies a broadly based multi-dimensional rehabilitative regimen. *West J Med*. 1974;120(5):386–391.
- Diamond L. Transgender experience and identity. In: Schwartz SJ, Luyckx K, Vignoles VL, eds. *Handbook of Identity Theory and Research*. New York, NY: Springer; 2011:629–647.
- Queen C, Schimmel L, eds. *PoMoSexuals: Challenging Assumptions About Gender and Sexuality*. San Francisco, CA: Cleis Press; 1997.
- Case LK, Ramachandran VS. Alternating gender incongruity: a new neuropsychiatric syndrome providing insight into the dynamic plasticity of brain sex. *Med Hypotheses*. 2012;78(5):626–631.
- Johnson TW, Wassersug RJ. Gender identity disorder outside the binary: when gender identity disorder not otherwise specified is not good enough. *Arch Sex Behav*. 2010;39(3):597–598.
- Wibowo E, Wassersug R, Warkentin K, Walker L, Robinson J, Brotto L, Johnson T. Impact of androgen deprivation therapy on sexual function: a response. *Asian J Androl*. 2012;14(5):793–794.
- Pasquosoone V. 7 countries giving transgender people fundamental rights the U.S. still won't. 2014. Available at: <https://mic.com/articles/87149/7-countries-giving-transgender-people-fundamental-rights-the-u-s-still-won-t>. Accessed 26 August 2016.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Arlington, VA: American Psychiatric Association Publishing.
- Drescher J, Cohen Kettner P, Winter S. Minding the body: situating gender identity diagnoses in the ICD 11. *Int Rev Psychiatry*. 2012;24(6):568–577.
- World Professional Association for Transgender Health. Standards of care for the health of transsexual, transgender, and gender nonconforming people. Available at: http://www.wpath.org/site/page.cfm?pk_association_webpage_menu_1351&pk_association_webpage_3926. Accessed 1 September 2017.
- Kreukels BP, Haraldsen IR, De Cuypere G, Richter Appelt H, Gijs L, Cohen Kettner PT. A European network for the investigation of gender incongruence: the ENIGI initiative. *Eur Psychiatry*. 2012;27(6):445–450.
- Dekker MJ, Wierckx K, Van Caenegem E, Klaver M, Kreukels BP, Elaut E, Fisher AD, van Trotsenburg MA, Schreiner T, den Heijer M, T'Sjoen G. A European network for the investigation of gender incongruence: endocrine part. *J Sex Med*. 2016;13(6):994–999.
- Ruble DN, Martin CL, Berenbaum SA. Gender development. In: Damon WL, Lerner RM, Eisenberg N, eds. *Handbook of Child Psychology: Social, Emotional, and Personality Development*. Vol. 3. 6th ed. New York, NY: Wiley; 2006:858–931.
- Steenma TD, Kreukels BP, de Vries AL, Cohen Kettner PT. Gender identity development in adolescence. *Horm Behav*. 2013;64(2):288–297.
- Rosenthal SM. Approach to the patient: transgender youth: endocrine considerations. *J Clin Endocrinol Metab*. 2014;99(12):4379–4389.
- Saraswat A, Weinand JD, Safer JD. Evidence supporting the biologic nature of gender identity. *Endocr Pract*. 2015;21(2):199–204.
- Gooren L. The biology of human psychosexual differentiation. *Horm Behav*. 2006;50(4):589–601.
- Berenbaum SA, Meyer Bahlburg HF. Gender development and sexuality in disorders of sex development. *Horm Metab Res*. 2015;47(5):361–366.
- Dessens AB, Slijper FME, Drop SLS. Gender dysphoria and gender change in chromosomal females with congenital adrenal hyperplasia. *Arch Sex Behav*. 2005;34(4):389–397.
- Meyer Bahlburg HFL, Dolezal C, Baker SW, Ehrhardt AA, New MI. Gender development in women with congenital adrenal hyperplasia as a function of disorder severity. *Arch Sex Behav*. 2006;35(6):667–684.
- Frisén L, Nordenstrom A, Falhammar H, Filipsson H, Holmdahl G, Janson PO, Thorén M, Hagenfeldt K, Moller A, Nordenskjöld A. Gender role behavior, sexuality, and psychosocial adaptation in women with congenital adrenal hyperplasia due to CYP21A2 deficiency. *J Clin Endocrinol Metab*. 2009;94(9):3432–3439.
- Meyer Bahlburg HFL, Dolezal C, Baker SW, Carlson AD, Obied JS, New MI. Prenatal androgenization affects gender related behavior but not gender identity in 5–12 year old girls with congenital adrenal hyperplasia. *Arch Sex Behav*. 2004;33(2):97–104.
- Cohen Kettner PT. Gender change in 46,XY persons with 5 α reductase 2 deficiency and 17 β hydroxysteroid dehydrogenase 3 deficiency. *Arch Sex Behav*. 2005;34(4):399–410.
- Reiner WG, Gearhart JP. Discordant sexual identity in some genetic males with cloacal exstrophy assigned to female sex at birth. *N Engl J Med*. 2004;350(4):333–341.
- Meyer Bahlburg HFL. Gender identity outcome in female raised 46,XY persons with penile agenesis, cloacal exstrophy of the bladder, or penile ablation. *Arch Sex Behav*. 2005;34(4):423–438.
- Coolidge FL, Thede LL, Young SE. The heritability of gender identity disorder in a child and adolescent twin sample. *Behav Genet*. 2002;32(4):251–257.
- Heylens G, De Cuypere G, Zucker KJ, Schelfaut C, Elaut E, Vanden Bossche H, De Baere E, T'Sjoen G. Gender identity disorder in twins: a review of the case report literature. *J Sex Med*. 2012;9(3):751–757.
- Fernández R, Esteva I, Gómez Gil E, Rumbo T, Almaraz MC, Roda E, Haro Mora J J, Guillamón A, Pávaro E. Association study of ER β , AR, and CYP19A1 genes and MtF transsexualism. *J Sex Med*. 2014;11(12):2986–2994.
- Henningsson S, Westberg L, Nilsson S, Lundstrom B, Ekselius L, Bodlund O, Lindstrom E, Hellstrand M, Rosmond R, Eriksson E, Landén M. Sex steroid related genes and male to female transsexualism. *Psychoneuroendocrinology*. 2005;30(7):657–664.
- Hare L, Bernard P, Sánchez FJ, Baird PN, Vilain E, Kennedy T, Harley VR. Androgen receptor repeat length polymorphism associated with male to female transsexualism. *Biol Psychiatry*. 2009;65(1):93–96.
- Lombardo F, Toselli L, Grassetti D, Paoli D, Masciandaro P, Valentini F, Lenzi A, Gandini L. Hormone and genetic study in

- male to female transsexual patients. *J Endocrinol Invest.* 2013; 36(8):550 557.
38. Ujike H, Otani K, Nakatsuka M, Ishii K, Sasaki A, Oishi T, Sato T, Okahisa Y, Matsumoto Y, Namba Y, Kimata Y, Kuroda S. As sociation study of gender identity disorder and sex hormone related genes. *Prog Neuropsychopharmacol Biol Psychiatry.* 2009;33(7):1241 1244.
 39. Kreukels BP, Guillamon A. Neuroimaging studies in people with gender incongruence. *Int Rev Psychiatry.* 2016;28(1): 120 128.
 40. Steensma TD, Biemond R, de Boer F, Cohen Kettenis PT. Desisting and persisting gender dysphoria after childhood: a qualitative follow up study. *Clin Child Psychol Psychiatry.* 2011;16(4):499 516.
 41. Wallien MSC, Cohen Kettenis PT. Psychosexual outcome of gender dysphoric children. *J Am Acad Child Adolesc Psychiatry.* 2008;47(12):1413 1423.
 42. Steensma TD, McGuire JK, Kreukels BPC, Beekman AJ, Cohen Kettenis PT. Factors associated with desistence and persistence of childhood gender dysphoria: a quantitative follow up study. *J Am Acad Child Adolesc Psychiatry.* 2013;52(6):582 590.
 43. Cohen Kettenis PT, Owen A, Kaijser VG, Bradley SJ, Zucker KJ. Demographic characteristics, social competence, and behavior problems in children with gender identity disorder: a cross national, cross clinic comparative analysis. *J Abnorm Child Psychol.* 2003;31(1):41 53.
 44. Dhejne C, Van Vlerken R, Heylens G, Arcelus J. Mental health and gender dysphoria: a review of the literature. *Int Rev Psychiatry.* 2016;28(1):44 57.
 45. Pasterski V, Gilligan L, Curtis R. Traits of autism spectrum dis orders in adults with gender dysphoria. *Arch Sex Behav.* 2014; 43(2):387 393.
 46. Spack NP, Edwards Leeper L, Feldman HA, Leibowitz S, Mandel F, Diamond DA, Vance SR. Children and adolescents with gender identity disorder referred to a pediatric medical center. *Pediatrics.* 2012;129(3):418 425.
 47. Terada S, Matsumoto Y, Sato T, Okabe N, Kishimoto Y, Uchitomi Y. Factors predicting psychiatric co morbidity in gender dysphoric adults. *Psychiatry Res.* 2012;200(2 3):469 474.
 48. VanderLaan DP, Leef JH, Wood H, Hughes SK, Zucker KJ. Autism spectrum disorder risk factors and autistic traits in gender dysphoric children. *J Autism Dev Disord.* 2015;45(6):1742 1750.
 49. de Vries ALC, Doreleijers TAH, Steensma TD, Cohen Kettenis PT. Psychiatric comorbidity in gender dysphoric adolescents. *J Child Psychol Psychiatry.* 2011;52(11):1195 1202.
 50. de Vries ALC, Noens ILJ, Cohen Kettenis PT, van Berckelaer Onnes IA, Doreleijers TA. Autism spectrum disorders in gender dysphoric children and adolescents. *J Autism Dev Disord.* 2010; 40(8):930 936.
 51. Wallien MSC, Swaab H, Cohen Kettenis PT. Psychiatric comorbidity among children with gender identity disorder. *J Am Acad Child Adolesc Psychiatry.* 2007;46(10):1307 1314.
 52. Kuiper AJ, Cohen Kettenis PT. Gender role reversal among postoperative transsexuals. Available at: https://www.atria.nl/ezines/web/IJT/97_03/numbers/symposium/ijtc0502.htm. Accessed 26 August 2016.
 53. Landén M, Wålinder J, Lambert G, Lundstrom B. Factors pre dictive of regret in sex reassignment. *Acta Psychiatr Scand.* 1998; 97(4):284 289.
 54. Olsson S E, Moller A. Regret after sex reassignment surgery in a male to female transsexual: a long term follow up. *Arch Sex Behav.* 2006;35(4):501 506.
 55. Pfafflin F, Junge A, eds. *Geschlechtsumwandlung: Abhand lungen zur Transsexualität.* Stuttgart, Germany: Schattauer; 1992.
 56. Lawrence AA. Factors associated with satisfaction or regret fol lowing male to female sex reassignment surgery. *Arch Sex Behav.* 2003;32(4):299 315.
 57. Cohen Kettenis PT, Pfafflin F. *Transgenderism and Intersexuality in Childhood and Adolescence: Making Choices.* Thousand Oaks, CA: SAGE Publications; 2003.
 58. Di Ceglie D, Freedman D, McPherson S, Richardson P. Children and adolescents referred to a specialist gender identity devel opment service: clinical features and demographic character istics. Available at: https://www.researchgate.net/publication/276061306_Children_and_Adolescents_Referred_to_a_Specialist_Gender_Identity_Development_Service_Clinical_Features_and_Demographic_Characteristics. Accessed 20 July 2017.
 59. Gijs L, Brewaeyns A. Surgical treatment of gender dysphoria in adults and adolescents: recent developments, effectiveness, and challenges. *Annu Rev Sex Res.* 2007;18:178 224.
 60. Cohen Kettenis PT, van Goozen SHM. Sex reassignment of ad olescent transsexuals: a follow up study. *J Am Acad Child Adolesc Psychiatry.* 1997;36(2):263 271.
 61. Smith YLS, van Goozen SHM, Cohen Kettenis PT. Adolescents with gender identity disorder who were accepted or rejected for sex reassignment surgery: a prospective follow up study. *J Am Acad Child Adolesc Psychiatry.* 2001;40(4):472 481.
 62. Smith YLS, Van Goozen SHM, Kuiper AJ, Cohen Kettenis PT. Sex reassignment: outcomes and predictors of treatment for adolescent and adult transsexuals. *Psychol Med.* 2005;35(1):89 99.
 63. de Vries ALC, McGuire JK, Steensma TD, Wagenaar ECF, Doreleijers TAH, Cohen Kettenis PT. Young adult psychological outcome after puberty suppression and gender reassignment. *Pediatrics.* 2014;134(4):696 704.
 64. Cole CM, O'Boyle M, Emory LE, Meyer WJ III. Comorbidity of gender dysphoria and other major psychiatric diagnoses. *Arch Sex Behav.* 1997;26(1):13 26.
 65. Cohen Kettenis PT, Schagen SEE, Steensma TD, de Vries ALC, Delemarre van de Waal HA. Puberty suppression in a gender dysphoric adolescent: a 22 year follow up. *Arch Sex Behav.* 2011; 40(4):843 847.
 66. First MB. Desire for amputation of a limb: paraphilia, psychosis, or a new type of identity disorder. *Psychol Med.* 2005;35(6): 919 928.
 67. Wierckx K, Van Caenegem E, Pennings G, Elaut E, Dedeker D, Van de Peer F, Weyers S, De Sutter P, T'Sjoen G. Reproductive wish in transsexual men. *Hum Reprod.* 2012;27(2):483 487.
 68. Wierckx K, Stuyver I, Weyers S, Hamada A, Agarwal A, De Sutter P, T'Sjoen G. Sperm freezing in transsexual women. *Arch Sex Behav.* 2012;41(5):1069 1071.
 69. Bertelloni S, Baroncelli GI, Ferdeghini M, Menchini Fabris F, Saggese G. Final height, gonadal function and bone mineral density of adolescent males with central precocious puberty after therapy with gonadotropin releasing hormone analogues. *Eur J Pediatr.* 2000;159(5):369 374.
 70. Buchter D, Behre HM, Kliesch S, Nieschlag E. Pulsatile GnRH or human chorionic gonadotropin/human menopausal gonadotro pin as effective treatment for men with hypogonadotropic hypogonadism: a review of 42 cases. *Eur J Endocrinol.* 1998; 139(3):298 303.
 71. Liu PY, Turner L, Rushford D, McDonald J, Baker HW, Conway AJ, Handelsman DJ. Efficacy and safety of recombinant human follicle stimulating hormone (Gonal F) with urinary human chorionic gonadotrophin for induction of spermatogenesis and fertility in gonadotrophin deficient men. *Hum Reprod.* 1999; 14(6):1540 1545.
 72. Pasquino AM, Pucarelli I, Accardo F, Demiraj V, Segni M, Di Nardo R. Long term observation of 87 girls with idiopathic central precocious puberty treated with gonadotropin releasing hormone analogs: impact on adult height, body mass index, bone mineral content, and reproductive function. *J Clin Endocrinol Metab.* 2008;93(1):190 195.
 73. Magiakou MA, Manousaki D, Papadaki M, Hadjidakis D, Levidou G, Vakaki M, Papaefstathiou A, Lalioti N, Kanaka Gantenbein C, Piaditis G, Chrousos GP, Dacou Voutetakis C. The

- efficacy and safety of gonadotropin releasing hormone analog treatment in childhood and adolescence: a single center, long term follow up study. *J Clin Endocrinol Metab.* 2010;**95**(1):109-117.
74. Baba T, Endo T, Honnma H, Kitajima Y, Hayashi T, Ikeda H, Masumori N, Kamiya H, Moriwaka O, Saito T. Association between polycystic ovary syndrome and female to male transsexuals. *Hum Reprod.* 2007;**22**(4):1011-1016.
 75. Spinder T, Spijkstra JJ, van den Tweel JG, Burger CW, van Kessel H, Hompes PGA, Gooren LJG. The effects of long term testosterone administration on pulsatile luteinizing hormone secretion and on ovarian histology in eugonadal female to male transsexual subjects. *J Clin Endocrinol Metab.* 1989;**69**(1):151-157.
 76. Baba T, Endo T, Ikeda K, Shimizu A, Honnma H, Ikeda H, Masumori N, Ohmura T, Kiya T, Fujimoto T, Koizumi M, Saito T. Distinctive features of female to male transsexualism and prevalence of gender identity disorder in Japan. *J Sex Med.* 2011;**8**(6):1686-1693.
 77. Vujovic S, Popovic S, Sbutega Milosevic G, Djordjevic M, Gooren L. Transsexualism in Serbia: a twenty year follow up study. *J Sex Med.* 2009;**6**(4):1018-1023.
 78. Ikeda K, Baba T, Noguchi H, Nagasawa K, Endo T, Kiya T, Saito T. Excessive androgen exposure in female to male transsexual persons of reproductive age induces hyperplasia of the ovarian cortex and stroma but not polycystic ovary morphology. *Hum Reprod.* 2013;**28**(2):453-461.
 79. Trebay G. He's pregnant. You're speechless. *New York Times.* 22 June 2008.
 80. Light AD, Obedin Maliver J, Sevelius JM, Kerns JL. Transgender men who experienced pregnancy after female to male gender transitioning. *Obstet Gynecol.* 2014;**124**(6):1120-1127.
 81. De Sutter P. Donor inseminations in partners of female to male transsexuals: should the question be asked? *Reprod Biomed Online.* 2003;**6**(3):382, author reply 282-283.
 82. De Roo C, Tilleman K, T'Sjoen G, De Sutter P. Fertility options in transgender people. *Int Rev Psychiatry.* 2016;**28**(1):112-119.
 83. Wennink JMB, Delemarre van de Waal HA, Schoemaker R, Schoemaker H, Schoemaker J. Luteinizing hormone and follicle stimulating hormone secretion patterns in boys throughout puberty measured using highly sensitive immunoradiometric assays. *Clin Endocrinol (Oxf).* 1989;**31**(5):551-564.
 84. Cohen Kettenis PT, Delemarre van de Waal HA, Gooren LJG. The treatment of adolescent transsexuals: changing insights. *J Sex Med.* 2008;**5**(8):1892-1897.
 85. Delemarre van de Waal HA, Cohen Kettenis PT. Clinical management of gender identity disorder in adolescents: a protocol on psychological and paediatric endocrinology aspects. *Eur J Endocrinol.* 2006;**155**:S131-S137.
 86. de Vries ALC, Steensma TD, Doreleijers TAH, Cohen Kettenis PT. Puberty suppression in adolescents with gender identity disorder: a prospective follow up study. *J Sex Med.* 2011;**8**(8):2276-2283.
 87. Bouman MB, van Zeijl MCT, Buncamper ME, Meijerink WJHJ, van Bodegraven AA, Muller MG. Intestinal vaginoplasty revisited: a review of surgical techniques, complications, and sexual function. *J Sex Med.* 2014;**11**(7):1835-1847.
 88. Carel JC, Eugster EA, Rogol A, Ghizzoni L, Palmert MR, Antoniazzi F, Berenbaum S, Bourguignon JP, Chrousos GP, Coste J, Deal S, de Vries L, Foster C, Heger S, Holland J, Jahnukainen K, Juul A, Kaplowitz P, Lahlou N, Lee MM, Lee P, Merke DP, Neely EK, Oostdijk W, Phillip M, Rosenfield RL, Shulman D, Styne D, Tauber M, Wit JM; ESPE LWPEES GnRH Analogs Consensus Conference Group. Consensus statement on the use of gonadotropin releasing hormone analogs in children. *Pediatrics.* 2009;**123**(4):e752-e762.
 89. Roth CL, Brendel L, Ruckert C, Hartmann K. Antagonistic and agonistic GnRH analogue treatment of precocious puberty: tracking gonadotropin concentrations in urine. *Horm Res.* 2005;**63**(5):257-262.
 90. Roth C. Therapeutic potential of GnRH antagonists in the treatment of precocious puberty. *Expert Opin Investig Drugs.* 2002;**11**(9):1253-1259.
 91. Tuvemo T. Treatment of central precocious puberty. *Expert Opin Investig Drugs.* 2006;**15**(5):495-505.
 92. Schagen SE, Cohen Kettenis PT, Delemarre van de Waal HA, Hannema SE. Efficacy and safety of gonadotropin releasing hormone agonist treatment to suppress puberty in gender dysphoric adolescents. *J Sex Med.* 2016;**13**(7):1125-1132.
 93. Manasco PK, Pescovitz OH, Feuillan PP, Hench KD, Barnes KM, Jones J, Hill SC, Loriaux DL, Cutler GB, Jr. Resumption of puberty after long term luteinizing hormone releasing hormone agonist treatment of central precocious puberty. *J Clin Endocrinol Metab.* 1988;**67**(2):368-372.
 94. Klink D, Caris M, Heijboer A, van Trotsenburg M, Rotteveel J. Bone mass in young adulthood following gonadotropin releasing hormone analog treatment and cross sex hormone treatment in adolescents with gender dysphoria. *J Clin Endocrinol Metab.* 2015;**100**(2):E270-E275.
 95. Finkelstein JS, Klibanski A, Neer RM. A longitudinal evaluation of bone mineral density in adult men with histories of delayed puberty. *J Clin Endocrinol Metab.* 1996;**81**(3):1152-1155.
 96. Bertelloni S, Baroncelli GI, Ferdeghini M, Perri G, Saggese G. Normal volumetric bone mineral density and bone turnover in young men with histories of constitutional delay of puberty. *J Clin Endocrinol Metab.* 1998;**83**(12):4280-4283.
 97. Darelid A, Ohlsson C, Nilsson M, Kindblom JM, Mellstrom D, Lorentzon M. Catch up in bone acquisition in young adult men with late normal puberty. *J Bone Miner Res.* 2012;**27**(10):2198-2207.
 98. Mittan D, Lee S, Miller E, Perez RC, Basler JW, Bruder JM. Bone loss following hypogonadism in men with prostate cancer treated with GnRH analogs. *J Clin Endocrinol Metab.* 2002;**87**(8):3656-3661.
 99. Saggese G, Bertelloni S, Baroncelli GI, Battini R, Franchi G. Reduction of bone density: an effect of gonadotropin releasing hormone analogue treatment in central precocious puberty. *Eur J Pediatr.* 1993;**152**(9):717-720.
 100. Neely EK, Bachrach LK, Hintz RL, Habiby RL, Slemenda CW, Feeze L, Pescovitz OH. Bone mineral density during treatment of central precocious puberty. *J Pediatr.* 1995;**127**(5):819-822.
 101. Bertelloni S, Baroncelli GI, Sorrentino MC, Perri G, Saggese G. Effect of central precocious puberty and gonadotropin releasing hormone analogue treatment on peak bone mass and final height in females. *Eur J Pediatr.* 1998;**157**(5):363-367.
 102. Thornton P, Silverman LA, Geffner ME, Neely EK, Gould E, Danoff TM. Review of outcomes after cessation of gonadotropin releasing hormone agonist treatment of girls with precocious puberty. *Pediatr Endocrinol Rev.* 2014;**11**(3):306-317.
 103. Lem AJ, van der Kaay DC, Hokken Koelega AC. Bone mineral density and body composition in short children born SGA during growth hormone and gonadotropin releasing hormone analog treatment. *J Clin Endocrinol Metab.* 2013;**98**(1):77-86.
 104. Antoniazzi F, Zamboni G, Bertoldo F, Lauriola S, Mengarda F, Pietrobelli A, Tatò L. Bone mass at final height in precocious puberty after gonadotropin releasing hormone agonist with and without calcium supplementation. *J Clin Endocrinol Metab.* 2003;**88**(3):1096-1101.
 105. Calcaterra V, Mannarino S, Corana G, Codazzi AC, Mazzola A, Brambilla P, Larizza D. Hypertension during therapy with triptorelin in a girl with precocious puberty. *Indian J Pediatr.* 2013;**80**(10):884-885.
 106. Siomou E, Kosmeri C, Pavlou M, Vlahos AP, Argyropoulou MI, Siamopoulou A. Arterial hypertension during treatment with triptorelin in a child with Williams Beuren syndrome. *Pediatr Nephrol.* 2014;**29**(9):1633-1636.
 107. Staphorsius AS, Kreukels BPC, Cohen Kettenis PT, Veltman DJ, Burke SM, Schagen SEE, Wouters FM, Delemarre van de Waal

- HA, Bakker J. Puberty suppression and executive functioning: an fMRI study in adolescents with gender dysphoria. *Psychoneuroendocrinology*. 2015;56:190–199.
108. Hough D, Bellingham M, Haraldsen IR, McLaughlin M, Rennie M, Robinson JE, Solbakk AK, Evans NP. Spatial memory is impaired by peripubertal GnRH agonist treatment and testosterone replacement in sheep. *Psychoneuroendocrinology*. 2017;75:173–182.
 109. Collipp PJ, Kaplan SA, Boyle DC, Plachte F, Kogut MD. Constitutional Isosexual Precocious Puberty. *Am J Dis Child*. 1964;108:399–405.
 110. Hahn HB, Jr, Hayles AB, Albert A. Medroxyprogesterone and constitutional precocious puberty. *Mayo Clin Proc*. 1964;39:182–190.
 111. Kaplan SA, Ling SM, Irani NG. Idiopathic isosexual precocity. *Am J Dis Child*. 1968;116(6):591–598.
 112. Schoen EJ. Treatment of idiopathic precocious puberty in boys. *J Clin Endocrinol Metab*. 1966;26(4):363–370.
 113. Gooren L. Hormone treatment of the adult transsexual patient. *Horm Res*. 2005;64(Suppl 2):31–36.
 114. Moore E, Wisniewski A, Dobs A. Endocrine treatment of transsexual people: a review of treatment regimens, outcomes, and adverse effects. *J Clin Endocrinol Metab*. 2003;88(8):3467–3473.
 115. Krueger RB, Hembree W, Hill M. Prescription of medroxyprogesterone acetate to a patient with pedophilia, resulting in Cushing's syndrome and adrenal insufficiency. *Sex Abuse*. 2006;18(2):227–228.
 116. Lynch MM, Khandheria MM, Meyer WJ. Retrospective study of the management of childhood and adolescent gender identity disorder using medroxyprogesterone acetate. *Int J Transgenderism*. 2015;16:201–208.
 117. Tack LJW, Craen M, Dhondt K, Vanden Bossche H, Laridaen J, Cools M. Consecutive lynestrenol and cross sex hormone treatment in biological female adolescents with gender dysphoria: a retrospective analysis. *Biol Sex Differ*. 2016;7:14.
 118. Hembree WC, Cohen Kettner P, Delemarre van de Waal HA, Gooren LJ, Meyer WJ 3rd, Spack NP, Tangpricha V, Montori VM; Endocrine Society. Endocrine treatment of transsexual persons: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2009;94(9):3132–3154.
 119. Mann L, Harmoni R, Power C. Adolescent decision making: the development of competence. *J Adolesc*. 1989;12(3):265–278.
 120. Stultiens L, Goffin T, Borry P, Dierickx K, Nys H. Minors and informed consent: a comparative approach. *Eur J Health Law*. 2007;14(1):21–46.
 121. Arshagouni P. "But I'm an adult now ... sort of". Adolescent consent in health care decision making and the adolescent brain. Available at: <http://digitalcommons.law.umaryland.edu/cgi/viewcontent.cgi?article=1124&context=jhclp>. Accessed 25 June 2017.
 122. NHS. Prescribing of cross sex hormones as part of the gender identity development service for children and adolescents. Available at: https://www.england.nhs.uk/commissioning/wp-content/uploads/sites/12/2016/08/clinical_com_pol_16046p.pdf. Accessed 14 June 2017.
 123. Ankarberg Lindgren C, Kristrom B, Norjavaara E. Physiological estrogen replacement therapy for puberty induction in girls: a clinical observational study. *Horm Res Paediatr*. 2014;81(4):239–244.
 124. Olson J, Schrager SM, Clark LF, Dunlap SL, Belzer M. Subcutaneous testosterone: an effective delivery mechanism for masculinizing young transgender men. *LGBT Health*. 2014;1(3):165–167.
 125. Spratt DI, Stewart I, Savage C, Craig W, Spack NP, Chandler DW, Spratt LV, Eimicke T, Olshan JS. Subcutaneous injection of testosterone is an effective and preferred alternative to intramuscular injection: demonstration in female to male transgender patients. *J Clin Endocrinol Metab*. 2017. doi:10.1210/jc.2017 00359
 126. Eisenegger C, von Eckardstein A, Fehr E, von Eckardstein S. Pharmacokinetics of testosterone and estradiol gel preparations in healthy young men. *Psychoneuroendocrinology*. 2013;38(2):171–178.
 127. de Ronde W, ten Kulve J, Woerdeman J, Kaufman J M, de Jong FH. Effects of oestradiol on gonadotrophin levels in normal and castrated men. *Clin Endocrinol (Oxf)*. 2009;71(6):874–879.
 128. Money J, Ehrhardt A. Man & woman, boy & girl: differentiation and dimorphism of gender identity from conception to maturity. Baltimore, MD: Johns Hopkins University Press; 1972:202–206.
 129. Heylens G, Verroken C, De Cock S, T'Sjoen G, De Cuypere G. Effects of different steps in gender reassignment therapy on psychopathology: a prospective study of persons with a gender identity disorder. *J Sex Med*. 2014;11(1):119–126.
 130. Costa R, Colizzi M. The effect of cross sex hormonal treatment on gender dysphoria individuals' mental health: a systematic review. *Neuropsychiatr Dis Treat*. 2016;12:1953–1966.
 131. Gooren LJG, Giltay EJ. Review of studies of androgen treatment of female to male transsexuals: effects and risks of administration of androgens to females. *J Sex Med*. 2008;5(4):765–776.
 132. Levy A, Crown A, Reid R. Endocrine intervention for transsexuals. *Clin Endocrinol (Oxf)*. 2003;59(4):409–418.
 133. Tangpricha V, Ducharme SH, Barber TW, Chipkin SR. Endocrinologic treatment of gender identity disorders. *Endocr Pract*. 2003;9(1):12–21.
 134. Meriggiola MC, Gava G. Endocrine care of transpeople part I. A review of cross sex hormonal treatments, outcomes and adverse effects in transmen. *Clin Endocrinol (Oxf)*. 2015;83(5):597–606.
 135. Bhasin S, Cunningham GR, Hayes FJ, Matsumoto AM, Snyder PJ, Swerdloff RS, Montori VM. Testosterone therapy in adult men with androgen deficiency syndromes: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab*. 2006;91(6):1995–2010.
 136. Pelusi C, Costantino A, Martelli V, Lambertini M, Bazzocchi A, Ponti F, Battista G, Venturoli S, Meriggiola MC. Effects of three different testosterone formulations in female to male transsexual persons. *J Sex Med*. 2014;11(12):3002–3011.
 137. Anderson GL, Limacher M, Assaf AR, Bassford T, Beresford SA, Clark H, Bonds D, Brunner R, Brzyski R, Caan B, Chlebowski R, Curb D, Gass M, Hays J, Heiss G, Hendrix S, Howard BV, Hsia J, Hubbell A, Jackson R, Johnson KC, Judd H, Kotchen JM, Kuller L, LaCroix AZ, Lane D, Langer RD, Lasser N, Lewis CE, Manson J, Margolis K, Ockene J, O'Sullivan MJ, Phillips L, Prentice RL, Ritenbaugh C, Robbins J, Rossouw JE, Sarto G, Stefanick ML, Van Horn L, Wactawski-Wende J, Wallace R, Wassertheil-Smoller S; Women's Health Initiative Steering Committee. Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: the Women's Health Initiative randomized controlled trial. *JAMA*. 2004;291(14):1701–1712.
 138. Dickersin K, Munro MG, Clark M, Langenberg P, Scherer R, Frick K, Zhu Q, Hallock L, Nichols J, Yalcinkaya TM; Surgical Treatments Outcomes Project for Dysfunctional Uterine Bleeding (STOP DUB) Research Group. Hysterectomy compared with endometrial ablation for dysfunctional uterine bleeding: a randomized controlled trial. *Obstet Gynecol*. 2007;110(6):1279–1289.
 139. Gooren LJ, Giltay EJ, Bunck MC. Long term treatment of transsexuals with cross sex hormones: extensive personal experience. *J Clin Endocrinol Metab*. 2008;93(1):19–25.
 140. Prior JC, Vigna YM, Watson D. Spironolactone with physiological female steroids for presurgical therapy of male to female transsexualism. *Arch Sex Behav*. 1989;18(1):49–57.
 141. Dittrich R, Binder H, Cupisti S, Hoffmann I, Beckmann MW, Mueller A. Endocrine treatment of male to female transsexuals using gonadotropin releasing hormone agonist. *Exp Clin Endocrinol Diabetes*. 2005;113(10):586–592.

142. Striipp B, Taylor AA, Bartter FC, Gillette JR, Loriaux DL, Easley R, Menard RH. Effect of spironolactone on sex hormones in man. *J Clin Endocrinol Metab.* 1975;41(4):777 781.
143. Levy J, Burshell A, Marbach M, Aflalo L, Glick SM. Interaction of spironolactone with oestradiol receptors in cytosol. *J Endocrinol.* 1980;84(3):371 379.
144. Wierckx K, Elaut E, Van Hoorde B, Heylens G, De Cuypere G, Monstrey S, Weyers S, Hoebeke P, T'Sjoen G. Sexual desire in trans persons: associations with sex reassignment treatment. *J Sex Med.* 2014;11(1):107 118.
145. Chiriaco G, Cauci S, Mazzon G, Trombetta C. An observational retrospective evaluation of 79 young men with long term adverse effects after use of finasteride against androgenetic alopecia. *Andrology.* 2016;4(2):245 250.
146. Gava G, Cerpolini S, Martelli V, Battista G, Seracchioli R, Meriggiola MC. Cyproterone acetate vs leuprolide acetate in combination with transdermal oestradiol in transwomen: a comparison of safety and effectiveness. *Clin Endocrinol (Oxf).* 2016; 85(2):239 246.
147. Casper RF, Yen SS. Rapid absorption of micronized estradiol 17 beta following sublingual administration. *Obstet Gynecol.* 1981; 57(1):62 64.
148. Price TM, Blauer KL, Hansen M, Stanczyk F, Lobo R, Bates GW. Single dose pharmacokinetics of sublingual versus oral administration of micronized 17 β estradiol. *Obstet Gynecol.* 1997;89(3): 340 345.
149. Toorians AWFT, Thomassen MCLGD, Zweegman S, Magdeleyns EJP, Tans G, Gooren LJG, Rosing J. Venous thrombosis and changes of hemostatic variables during cross sex hormone treatment in transsexual people. *J Clin Endocrinol Metab.* 2003;88(12): 5723 5729.
150. Mepham N, Bouman WP, Arcelus J, Hayter M, Wylie KR. People with gender dysphoria who self prescribe cross sex hormones: prevalence, sources, and side effects knowledge. *J Sex Med.* 2014; 11(12):2995 3001.
151. Richards C, Bouman WP, Seal L, Barker MJ, Nieder TO, T'Sjoen G. Non binary or genderqueer genders. *Int Rev Psychiatry.* 2016; 28(1):95 102.
152. Cosyns M, Van Borsel J, Wierckx K, Dedeker D, Van de Peer F, Daelman T, Laenen S, T'Sjoen G. Voice in female to male transsexual persons after long term androgen therapy. *Laryngoscope.* 2014;124(6):1409 1414.
153. Deuster D, Matulat P, Knief A, Zitzmann M, Rosslau K, Szukaj M, am Zehnhoff Dinnesen A, Schmidt CM. Voice deepening under testosterone treatment in female to male gender dysphoric individuals. *Eur Arch Otorhinolaryngol.* 2016;273(4):959 965.
154. Lapauw B, Taes Y, Simoens S, Van Caenegem E, Weyers S, Goemaere S, Toye K, Kaufman J M, T'Sjoen GG. Body composition, volumetric and areal bone parameters in male to female transsexual persons. *Bone.* 2008;43(6):1016 1021.
155. Meyer III WJ, Webb A, Stuart CA, Finkelstein JW, Lawrence B, Walker PA. Physical and hormonal evaluation of transsexual patients: a longitudinal study. *Arch Sex Behav.* 1986;15(2): 121 138.
156. Asscheman H, Gooren LJ, Assies J, Smits JP, de Slegte R. Prolactin levels and pituitary enlargement in hormone treated male to female transsexuals. *Clin Endocrinol (Oxf).* 1988;28(6):583 588.
157. Gooren LJ, Harmsen Louman W, van Kessel H. Follow up of prolactin levels in long term oestrogen treated male to female transsexuals with regard to prolactinoma induction. *Clin Endocrinol (Oxf).* 1985;22(2):201 207.
158. Wierckx K, Van Caenegem E, Schreiner T, Haraldsen I, Fisher AD, Toye K, Kaufman JM, T'Sjoen G. Cross sex hormone therapy in trans persons is safe and effective at short time follow up: results from the European network for the investigation of gender incongruence. *J Sex Med.* 2014;11(8):1999 2011.
159. Ott J, Kaufmann U, Bentz EK, Huber JC, Tempfer CB. Incidence of thrombophilia and venous thrombosis in transsexuals under cross sex hormone therapy. *Fertil Steril.* 2010;93(4):1267 1272.
160. Giltay EJ, Hoogeveen EK, Elbers JMH, Gooren LJG, Asscheman H, Stehouwer CDA. Effects of sex steroids on plasma total homocysteine levels: a study in transsexual males and females. *J Clin Endocrinol Metab.* 1998;83(2):550 553.
161. van Kesteren PJM, Asscheman H, Megens JAJ, Gooren LJG. Mortality and morbidity in transsexual subjects treated with cross sex hormones. *Clin Endocrinol (Oxf).* 1997;47(3): 337 343.
162. Wierckx K, Gooren L, T'Sjoen G. Clinical review: breast development in trans women receiving cross sex hormones. *J Sex Med.* 2014;11(5):1240 1247.
163. Bird D, Vowles K, Anthony PP. Spontaneous rupture of a liver cell adenoma after long term methyltestosterone: report of a case successfully treated by emergency right hepatic lobectomy. *Br J Surg.* 1979;66(3):212 213.
164. Westaby D, Ogle SJ, Paradinas FJ, Randell JB, Murray Lyon IM. Liver damage from long term methyltestosterone. *Lancet.* 1977; 2(8032):262 263.
165. Weinand JD, Safer JD. Hormone therapy in transgender adults is safe with provider supervision; a review of hormone therapy sequelae for transgender individuals. *J Clin Transl Endocrinol.* 2015;2(2):55 60.
166. Roberts TK, Kraft CS, French D, Ji W, Wu AH, Tangpricha V, Fantz CR. Interpreting laboratory results in transgender patients on hormone therapy. *Am J Med.* 2014;127(2):159 162.
167. Vesper HW, Botelho JC, Wang Y. Challenges and improvements in testosterone and estradiol testing. *Asian J Androl.* 2014;16(2): 178 184.
168. Asscheman H, T'Sjoen G, Lemaire A, Mas M, Meriggiola MC, Mueller A, Kuhn A, Dhejne C, Morel Journal N, Gooren LJ. Venous thrombo embolism as a complication of cross sex hormone treatment of male to female transsexual subjects: a review. *Andrologia.* 2014;46(7):791 795.
169. Righini M, Perrier A, De Moerloose P, Bounameaux H. D dimer for venous thromboembolism diagnosis: 20 years later. *J Thromb Haemost.* 2008;6(7):1059 1071.
170. Gooren LJ, Assies J, Asscheman H, de Slegte R, van Kessel H. Estrogen induced prolactinoma in a man. *J Clin Endocrinol Metab.* 1988;66(2):444 446.
171. Kovacs K, Stefaneanu L, Ezzat S, Smyth HS. Prolactin producing pituitary adenoma in a male to female transsexual patient with protracted estrogen administration. A morphologic study. *Arch Pathol Lab Med.* 1994;118(5):562 565.
172. Serri O, Noiseux D, Robert F, Hardy J. Lactotroph hyperplasia in an estrogen treated male to female transsexual patient. *J Clin Endocrinol Metab.* 1996;81(9):3177 3179.
173. Cunha FS, Domenice S, Câmara VL, Sircili MH, Gooren LJ, Mendonça BB, Costa EM. Diagnosis of prolactinoma in two male to female transsexual subjects following high dose cross sex hormone therapy. *Andrologia.* 2015;47(6):680 684.
174. Nota NM, Dekker MJHJ, Klaver M, Wiepjes CM, van Trotsenburg MA, Heijboer AC, den Heijer M. Prolactin levels during short and long term cross sex hormone treatment: an observational study in transgender persons. *Andrologia.* 2017;49(6).
175. Bunck MC, Debono M, Giltay EJ, Verheijen AT, Diamant M, Gooren LJ. Autonomous prolactin secretion in two male to female transgender patients using conventional oestrogen doses. *BMJ Case Rep.* 2009;2009:bcr0220091589.
176. Elamin MB, Garcia MZ, Murad MH, Erwin PJ, Montori VM. Effect of sex steroid use on cardiovascular risk in transsexual individuals: a systematic review and meta analyses. *Clin Endocrinol (Oxf).* 2010;72(1):1 10.
177. Berra M, Armillotta F, D'Emidio L, Costantino A, Martorana G, Pelusi G, Meriggiola MC. Testosterone decreases adiponectin

- levels in female to male transsexuals. *Asian J Androl.* 2006;8(6):725 729.
178. Elbers JMH, Giltay EJ, Teerlink T, Scheffer PG, Asscheman H, Seidell JC, Gooren LJG. Effects of sex steroids on components of the insulin resistance syndrome in transsexual subjects. *Clin Endocrinol (Oxf).* 2003;58(5):562 571.
 179. Giltay EJ, Lambert J, Gooren LJG, Elbers JMH, Steyn M, Stehouwer CDA. Sex steroids, insulin, and arterial stiffness in women and men. *Hypertension.* 1999;34(4 Pt 1):590 597.
 180. Polderman KH, Gooren LJ, Asscheman H, Bakker A, Heine RJ. Induction of insulin resistance by androgens and estrogens. *J Clin Endocrinol Metab.* 1994;79(1):265 271.
 181. Maraka S. Effect of sex steroids on lipids, venous thromboembolism, cardiovascular disease and mortality in transgender individuals: a systematic review and meta analysis. Available at: <http://press.endocrine.org/doi/abs/10.1210/endo-meetings.2016.RE.15.FRI.136>. Accessed 3 July 2017.
 182. Meriggiola MC, Armillotta F, Costantino A, Altieri P, Saad F, Kalhorn T, Perrone AM, Ghi T, Pelusi C, Pelusi G. Effects of testosterone undecanoate administered alone or in combination with letrozole or dutasteride in female to male transsexuals. *J Sex Med.* 2008;5(10):2442 2453.
 183. Giltay EJ, Toorians AW, Sarabdjitsingh AR, de Vries NA, Gooren LJ. Established risk factors for coronary heart disease are unrelated to androgen induced baldness in female to male transsexuals. *J Endocrinol.* 2004;180(1):107 112.
 184. Giltay EJ, Verhoef P, Gooren LJG, Geleijnse JM, Schouten EG, Stehouwer CDA. Oral and transdermal estrogens both lower plasma total homocysteine in male to female transsexuals. *Ath erosclerosis.* 2003;168(1):139 146.
 185. Calof OM, Singh AB, Lee ML, Kenny AM, Urban RJ, Tenover JL, Bhasin S. Adverse events associated with testosterone replacement in middle aged and older men: a meta analysis of randomized, placebo controlled trials. *J Gerontol A Biol Sci Med Sci.* 2005;60(11):1451 1457.
 186. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA.* 2001;285(19):2486 2497.
 187. Murad MH, Elamin MB, Garcia MZ, Mullan RJ, Murad A, Erwin PJ, Montori VM. Hormonal therapy and sex reassignment: a systematic review and meta analysis of quality of life and psychosocial outcomes. *Clin Endocrinol (Oxf).* 2010;72(2):214 231.
 188. Van Caenegem E, Wierckx K, Taes Y, Schreiner T, Vandewalle S, Toye K, Lapauw B, Kaufman JM, T'Sjoen G. Body composition, bone turnover, and bone mass in trans men during testosterone treatment: 1 year follow up data from a prospective case controlled study (ENIGI). *Eur J Endocrinol.* 2015;172(2):163 171.
 189. Turner A, Chen TC, Barber TW, Malabanan AO, Holick MF, Tangpricha V. Testosterone increases bone mineral density in female to male transsexuals: a case series of 15 subjects. *Clin Endocrinol (Oxf).* 2004;61(5):560 566.
 190. van Kesteren P, Lips P, Gooren LJG, Asscheman H, Megens J. Long term follow up of bone mineral density and bone metabolism in transsexuals treated with cross sex hormones. *Clin Endocrinol (Oxf).* 1998;48(3):347 354.
 191. Van Caenegem E, Taes Y, Wierckx K, Vandewalle S, Toye K, Kaufman JM, Schreiner T, Haraldsen I, T'Sjoen G. Low bone mass is prevalent in male to female transsexual persons before the start of cross sex hormonal therapy and gonadectomy. *Bone.* 2013;54(1):92 97.
 192. Amin S, Zhang Y, Sawin CT, Evans SR, Hannan MT, Kiel DP, Wilson PW, Felson DT. Association of hypogonadism and estradiol levels with bone mineral density in elderly men from the Framingham study. *Ann Intern Med.* 2000;133(12):951 963.
 193. Gennari L, Khosla S, Bilezikian JP. Estrogen and fracture risk in men. *J Bone Miner Res.* 2008;23(10):1548 1551.
 194. Khosla S, Melton LJ III, Atkinson EJ, O'Fallon WM, Klee GG, Riggs BL. Relationship of serum sex steroid levels and bone turnover markers with bone mineral density in men and women: a key role for bioavailable estrogen. *J Clin Endocrinol Metab.* 1998;83(7):2266 2274.
 195. Mueller A, Dittrich R, Binder H, Kuehnel W, Maltaris T, Hoffmann I, Beckmann MW. High dose estrogen treatment increases bone mineral density in male to female transsexuals receiving gonadotropin releasing hormone agonist in the absence of testosterone. *Eur J Endocrinol.* 2005;153(1):107 113.
 196. Ruetsche AG, Kneubuehl R, Birkhaeuser MH, Lippuner K. Cortical and trabecular bone mineral density in transsexuals after long term cross sex hormonal treatment: a cross sectional study. *Osteoporos Int.* 2005;16(7):791 798.
 197. Ganly I, Taylor EW. Breast cancer in a trans sexual man receiving hormone replacement therapy. *Br J Surg.* 1995;82(3):341.
 198. Pritchard TJ, Pankowsky DA, Crowe JP, Abdul Karim FW. Breast cancer in a male to female transsexual. A case report. *JAMA.* 1988;259(15):2278 2280.
 199. Symmers WS. Carcinoma of breast in trans sexual individuals after surgical and hormonal interference with the primary and secondary sex characteristics. *BMJ.* 1968;2(5597):83 85.
 200. Brown GR. Breast cancer in transgender veterans: a ten case series. *LGBT Health.* 2015;2(1):77 80.
 201. Shao T, Grossbard ML, Klein P. Breast cancer in female to male transsexuals: two cases with a review of physiology and management. *Clin Breast Cancer.* 2011;11(6):417 419.
 202. Nikolic DV, Djordjevic ML, Granic M, Nikolic AT, Stanimirovic VV, Zdravkovic D, Jelic S. Importance of revealing a rare case of breast cancer in a female to male transsexual after bilateral mastectomy. *World J Surg Oncol.* 2012;10:280.
 203. Bosze P, Tóth A, Torok M. Hormone replacement and the risk of breast cancer in Turner's syndrome. *N Engl J Med.* 2006;355(24):2599 2600.
 204. Schoemaker MJ, Swerdlow AJ, Higgins CD, Wright AF, Jacobs PA; UK Clinical Cytogenetics Group. Cancer incidence in women with Turner syndrome in Great Britain: a national cohort study. *Lancet Oncol.* 2008;9(3):239 246.
 205. Smith RA, Cokkinides V, Eyre HJ. American Cancer Society guidelines for the early detection of cancer, 2006. *CA Cancer J Clin.* 2006;56(1):11 25, quiz 49 50.
 206. Wilson JD, Roehrborn C. Long term consequences of castration in men: lessons from the Skoptzy and the eunuchs of the Chinese and Ottoman courts. *J Clin Endocrinol Metab.* 1999;84(12):4324 4331.
 207. van Kesteren P, Meinhardt W, van der Valk P, Geldof A, Megens J, Gooren L. Effects of estrogens only on the prostates of aging men. *J Urol.* 1996;156(4):1349 1353.
 208. Brown JA, Wilson TM. Benign prostatic hyperplasia requiring transurethral resection of the prostate in a 60 year old male to female transsexual. *Br J Urol.* 1997;80(6):956 957.
 209. Casella R, Bubendorf L, Schaefer DJ, Bachmann A, Gasser TC, Sulser T. Does the prostate really need androgens to grow? Transurethral resection of the prostate in a male to female transsexual 25 years after sex changing operation. *Urol Int.* 2005;75(3):288 290.
 210. Dorff TB, Shazer RL, Nepomuceno EM, Tucker SJ. Successful treatment of metastatic androgen independent prostate carcinoma in a transsexual patient. *Clin Genitourin Cancer.* 2007;5(5):344 346.
 211. Thurston AV. Carcinoma of the prostate in a transsexual. *Br J Urol.* 1994;73(2):217.

212. van Harst EP, Newling DW, Gooren LJ, Asscheman H, Prenger DM. Metastatic prostatic carcinoma in a male to female transsexual. *BJU Int*. 1998;81:776.
213. Turo R, Jallad S, Prescott S, Cross WR. Metastatic prostate cancer in transsexual diagnosed after three decades of estrogen therapy. *Can Urol Assoc J*. 2013;7(7-8):E544 E546.
214. Miksad RA, Bubley G, Church P, Sanda M, Rofsky N, Kaplan I, Cooper A. Prostate cancer in a transgender woman 41 years after initiation of feminization. *JAMA*. 2006;296(19):2316 2317.
215. Moyer VA; U.S. Preventive Services Task Force. Screening for prostate cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2012;157(2):120 134.
216. Futterweit W. Endocrine therapy of transsexualism and potential complications of long term treatment. *Arch Sex Behav*. 1998;27(2):209 226.
217. Miller N, Bédard YC, Cooter NB, Shaul DL. Histological changes in the genital tract in transsexual women following androgen therapy. *Histopathology*. 1986;10(7):661 669.
218. O'Hanlan KA, Dibble SL, Young Spint M. Total laparoscopic hysterectomy for female to male transsexuals. *Obstet Gynecol*. 2007;110(5):1096 1101.
219. Dizon DS, Tejada Berges T, Koelliker S, Steinhoff M, Granai CO. Ovarian cancer associated with testosterone supplementation in a female to male transsexual patient. *Gynecol Obstet Invest*. 2006;62(4):226 228.
220. Hage JJ, Dekker JJML, Karim RB, Verheijen RHM, Bloemena E. Ovarian cancer in female to male transsexuals: report of two cases. *Gynecol Oncol*. 2000;76(3):413 415.
221. Mueller A, Gooren L. Hormone related tumors in transsexuals receiving treatment with cross sex hormones. *Eur J Endocrinol*. 2008;159(3):197 202.
222. Coleman E, Bockting W, Botzer M, Cohen Kettner P, DeCuypere G, Feldman J, Fraser L, Green J, Knudson G, Meyer WJ, Monstrey S, Adler RK, Brown GR, Devor AH, Ehrbar R, Ettner R, Eyster E, Garofalo R, Karasic DH, Lev AL, Mayer G, Meyer Bahlburg H, Hall BP, Pfaefflin F, Rachlin K, Robinson B, Schechter LS, Tangpricha V, van Trotsenburg M, Vitale A, Winter S, Whittle S, Wylie KR, Zucker K. Standards of care for the health of transsexual, transgender, and gender nonconforming people, version 7. *Int J Transgenderism*. 2012;13:165 232.
223. Colebunders B, D'Arpa S, Weijers S, Lumen N, Hoebeke P, Monstrey S. Female to male gender reassignment surgery. In: Ettner R, Monstrey S, Coleman E, eds. *Principles of Transgender Medicine and Surgery*. 2nd ed. New York, NY: Routledge Taylor & Francis Group; 2016:279 317.
224. Monstrey S, Hoebeke P, Dhont M, De Cuypere G, Rubens R, Moerman M, Hamdi M, Van Landuyt K, Blondeel P. Surgical therapy in transsexual patients: a multi disciplinary approach. *Acta Chir Belg*. 2001;101(5):200 209.
225. Selvaggi G, Ceulemans P, De Cuypere G, VanLanduyt K, Blondeel P, Hamdi M, Bowman C, Monstrey S. Gender identity disorder: general overview and surgical treatment for vaginoplasty in male to female transsexuals. *Plast Reconstr Surg*. 2005;116(6):135e 145e.
226. Tugnet N, Goddard JC, Vickery RM, Khoosal D, Terry TR. Current management of male to female gender identity disorder in the UK. *Postgrad Med J*. 2007;83(984):638 642.
227. Horbach SER, Bouman M B, Smit JM, Ozer M, Buncamper ME, Mullender MG. Outcome of vaginoplasty in male to female transsexuals: a systematic review of surgical techniques. *J Sex Med*. 2015;12(6):1499 1512.
228. Wroblewski P, Gustafsson J, Selvaggi G. Sex reassignment surgery for transsexuals. *Curr Opin Endocrinol Diabetes Obes*. 2013;20(6):570 574.
229. Morrison SD, Satterwhite T, Grant DW, Kirby J, Laub DR, Sr, VanMaasdam J. Long term outcomes of rectosigmoid neo colporrhaphy in male to female gender reassignment surgery. *Plast Reconstr Surg*. 2015;136(2):386 394.
230. Dessy LA, Mazzocchi M, Corrias F, Ceccarelli S, Marchese C, Scuderi N. The use of cultured autologous oral epithelial cells for vaginoplasty in male to female transsexuals: a feasibility, safety, and advantageousness clinical pilot study. *Plast Reconstr Surg*. 2014;133(1):158 161.
231. Li FY, Xu YS, Zhou CD, Zhou Y, Li SK, Li Q. Long term outcomes of vaginoplasty with autologous buccal micromucosa. *Obstet Gynecol*. 2014;123(5):951 956.
232. Kanhai RC. Sensate vagina pedicled spot for male to female transsexuals: the experience in the first 50 patients. *Aesthetic Plast Surg*. 2016;40(2):284 287.
233. Straayer C. Transplants for transsexuals? Ambitions, concerns, ideology. Paper presented at: Trans*Studies: An International Transdisciplinary Conference on Gender, Embodiment, and Sexuality; 7-10 September 2016; University of Arizona, Tucson, AZ.
234. Bucci S, Mazzon G, Liguori G, Napoli R, Pavan N, Bormioli S, Olandini G, De Concilio B, Trombetta C. Neovaginal prolapse in male to female transsexuals: an 18 year long experience. *Biomed Res Int*. 2014;2014:240761.
235. Raigosa M, Avvedimento S, Yoon TS, Cruz Gimeno J, Rodriguez G, Fontdevila J. Male to female genital reassignment surgery: a retrospective review of surgical technique and complications in 60 patients. *J Sex Med*. 2015;12(8):1837 1845.
236. Green R. Sexual functioning in post operative transsexuals: male to female and female to male. *Int J Impot Res*. 1998;10(Suppl 1):S22 S24.
237. Hess J, Rossi Neto R, Panic L, Rubben H, Senf W. Satisfaction with male to female gender reassignment surgery. *Dtsch Arztebl Int*. 2014;111(47):795 801.
238. Nygren U, Nordenskjold A, Arver S, Sodersten M. Effects on voice fundamental frequency and satisfaction with voice in trans men during testosterone treatment a longitudinal study. *J Voice*. 2016;30(6):766.e23 766.e34.
239. Becking AG, Tuinzing DB, Hage JJ, Gooren LJG. Transgender feminization of the facial skeleton. *Clin Plast Surg*. 2007;34(3):557 564.
240. Giraldo F, Esteva I, Bergero T, Cano G, González C, Salinas P, Rivada E, Lara JS, Soriguer F; Andalusia Gender Team. Corona glans clitoroplasty and urethrapreputial vestibuloplasty in male to female transsexuals: the vulval aesthetic refinement by the Andalusia Gender Team. *Plast Reconstr Surg*. 2004;114(6):1543 1550.
241. Goddard JC, Vickery RM, Terry TR. Development of feminizing genitoplasty for gender dysphoria. *J Sex Med*. 2007;4(4 Pt 1):981 989.
242. Hage JJ, de Graaf FH, Bouman FG, Bloem JJAM. Sculpturing the glans in phalloplasty. *Plast Reconstr Surg*. 1993;92(1):157 161, discussion 162.
243. Thiagaraj D, Gunasegaram R, Loganath A, Peh KL, Kottegoda SR, Ratnam SS. Histopathology of the testes from male transsexuals on oestrogen therapy. *Ann Acad Med Singapore*. 1987;16(2):347 348.
244. Monstrey SJ, Ceulemans P, Hoebeke P. Sex reassignment surgery in the female to male transsexual. *Semin Plast Surg*. 2011;25(3):229 244.
245. Perovic SV, Djinovic R, Bumbasirevic M, Djordjevic M, Vukovic P. Total phalloplasty using a musculocutaneous latissimus dorsi flap. *BJU Int*. 2007;100(4):899 905, discussion 905.
246. Vesely J, Hyza P, Ranno R, Cigna E, Monni N, Stupka I, Justan I, Dvorak Z, Novak P, Ranno S. New technique of total phalloplasty with reinnervated latissimus dorsi myocutaneous free flap in female to male transsexuals. *Ann Plast Surg*. 2007;58(5):544 550.
247. Ranno R, Vesely J, Hýza P, Stupka I, Justan I, Dvorák Z, Monni N, Novák P, Ranno S. Neo phalloplasty with re innervated latissimus dorsi free flap: a functional study of a novel technique. *Acta Chir Plast*. 2007;49(1):3 7.

248. Garcia MM, Christopher NA, De Luca F, Spilotros M, Ralph DJ. Overall satisfaction, sexual function, and the durability of neophallus dimensions following staged female to male genital gender confirming surgery: the Institute of Urology, London U.K. experience. *Transl Androl Urol.* 2014;3(2):156 162.
249. Chen H C, Gedebou TM, Yazar S, Tang Y B. Prefabrication of the free fibula osteocutaneous flap to create a functional human penis using a controlled fistula method. *J Reconstr Microsurg.* 2007;23(3):151 154.
250. Hoebeke PB, Decaestecker K, Beysens M, Opdenakker Y, Lumen N, Monstrey SM. Erectile implants in female to male transsexuals: our experience in 129 patients. *Eur Urol.* 2010;57(2):334 341.
251. Hage JJ. Metaidoioplasty: an alternative phalloplasty technique in transsexuals. *Plast Reconstr Surg.* 1996;97(1):161 167.
252. Cohanad S. Extensive metoidioplasty as a technique capable of creating a compatible analogue to a natural penis in female transsexuals. *Aesthetic Plast Surg.* 2016;40(1):130 138.
253. Selvaggi G, Hoebeke P, Ceulemans P, Hamdi M, Van Landuyt K, Blondeel P, De Cuyper G, Monstrey S. Scrotal reconstruction in female to male transsexuals: a novel scrotoplasty. *Plast Reconstr Surg.* 2009;123(6):1710 1718.
254. Bjerrome Ahlin H, Kolby L, Elander A, Selvaggi G. Improved results after implementation of the Ghent algorithm for subcutaneous mastectomy in female to male transsexuals. *J Plast Surg Hand Surg.* 2014;48(6):362 367.
255. Wolter A, Diedrichson J, Scholz T, Arens Landwehr A, Liebau J. Sexual reassignment surgery in female to male transsexuals: an algorithm for subcutaneous mastectomy. *J Plast Reconstr Aesthet Surg.* 2015;68(2):184 191.
256. Richards C, Barrett J. The case for bilateral mastectomy and male chest contouring for the female to male transsexual. *Ann R Coll Surg Engl.* 2013;95(2):93 95.
257. Sutcliffe PA, Dixon S, Akehurst RL, Wilkinson A, Shippam A, White S, Richards R, Caddy CM. Evaluation of surgical procedures for sex reassignment: a systematic review. *J Plast Reconstr Aesthet Surg.* 2009;62(3):294 306, discussion 306 308.
258. Selvaggi G, Elander A. Penile reconstruction/formation. *Curr Opin Urol.* 2008;18(6):589 597.
259. Dhejne C, Lichtenstein P, Boman M, Johansson ALV, Långström N, Landén M. Long term follow up of transsexual persons undergoing sex reassignment surgery: cohort study in Sweden. *PLoS One.* 2011;6(2):e16885.
260. Kuhn A, Bodmer C, Stadlmayr W, Kuhn P, Mueller MD, Birkhauser M. Quality of life 15 years after sex reassignment surgery for transsexualism. *Fertil Steril.* 2009;92(5):1685 1689.e3.
261. Papadopoulos NA, Lellé JD, Zavlin D, Herschbach P, Henrich G, Kovacs L, Ehrenberger B, Kluger AK, Machens HG, Schaff J. Quality of life and patient satisfaction following male to female sex reassignment surgery. *J Sex Med.* 2017;14(5):721 730.
262. Simonsen RK, Hald GM, Kristensen E, Giraldo A. Long term follow up of individuals undergoing sex reassignment surgery: somatic morbidity and cause of death. *Sex Med.* 2016;4(1):e60 e68.
263. Djordjevic ML, Bizic MR, Duisin D, Bouman MB, Buncamper M. Reversal Surgery in regretful male to female transsexuals after sex reassignment surgery. *J Sex Med.* 2016;13(6):1000 1007.
264. Liberopoulos EN, Florentin M, Mikhailidis DP, Elisaf MS. Compliance with lipid lowering therapy and its impact on cardiovascular morbidity and mortality. *Expert Opin Drug Saf.* 2008;7(6):717 725.
265. Forbes SS, Stephen WJ, Harper WL, Loeb M, Smith R, Christoffersen EP, McLean RF. Implementation of evidence based practices for surgical site infection prophylaxis: results of a pre and postintervention study. *J Am Coll Surg.* 2008;207(3):336 341.
266. Davis PJ, Spady D, de Gara C, Forgie SE. Practices and attitudes of surgeons toward the prevention of surgical site infections: a provincial survey in Alberta, Canada. *Infect Control Hosp Epidemiol.* 2008;29(12):1164 1166.

EXHIBIT 19



Standards of Care for the Health of Transsexual, Transgender, and Gender Nonconforming People

The World Professional Association for Transgender Health



Standards of Care for the Health of Transsexual, Transgender, and Gender Nonconforming People

The World Professional Association for Transgender Health

7th Version¹ | www.wpath.org

¹ This is the seventh version of the Standards of Care. The original SOC were published in 1979. Previous revisions were in 1980, 1981, 1990, 1998, and 2001.

Table of Contents

I. Purpose and Use of the Standards of Care	1
II. Global Applicability of the Standards of Care	3
III. The Difference between Gender Nonconformity and Gender Dysphoria	4
IV. Epidemiologic Considerations	6
V. Overview of Therapeutic Approaches for Gender Dysphoria	8
VI. Assessment and Treatment of Children and Adolescents with Gender Dysphoria	10
VII. Mental Health	21
VIII. Hormone Therapy	33
IX. Reproductive Health	50
X. Voice and Communication Therapy	52
XI. Surgery	54
XII. Postoperative Care and Follow-Up	64
XIII. Lifelong Preventive and Primary Care	65
XIV. Applicability of the <i>Standards of Care</i> to People Living in Institutional Environments	67
XV. Applicability of the <i>Standards of Care</i> to People With Disorders of Sex Development	69
References	72
Appendices:	
A. Glossary	95
B. Overview of Medical Risks of Hormone Therapy	97
C. Summary of Criteria for Hormone Therapy and Surgeries	104
D. Evidence for Clinical Outcomes of Therapeutic Approaches	107
E. Development Process for the <i>Standards of Care, Version 7</i>	109

- Surgery to change primary and/or secondary sex characteristics (e.g., breasts/chest, external and/or internal genitalia, facial features, body contouring);
- Psychotherapy (individual, couple, family, or group) for purposes such as exploring gender identity, role, and expression; addressing the negative impact of gender dysphoria and stigma on mental health; alleviating internalized transphobia; enhancing social and peer support; improving body image; or promoting resilience.

Options for Social Support and Changes in Gender Expression

In addition (or as an alternative) to the psychological and medical treatment options described above, other options can be considered to help alleviate gender dysphoria, for example:

- Offline and online peer support resources, groups, or community organizations that provide avenues for social support and advocacy;
- Offline and online support resources for families and friends;
- Voice and communication therapy to help individuals develop verbal and non-verbal communication skills that facilitate comfort with their gender identity;
- Hair removal through electrolysis, laser treatment, or waxing;
- Breast binding or padding, genital tucking or penile prostheses, padding of hips or buttocks;
- Changes in name and gender marker on identity documents.

VI

Assessment and Treatment of Children and Adolescents with Gender Dysphoria

There are a number of differences in the phenomenology, developmental course, and treatment approaches for gender dysphoria in children, adolescents, and adults. In children and adolescents, a rapid and dramatic developmental process (physical, psychological, and sexual) is involved and

there is greater fluidity and variability in outcomes, particular in prepubertal children. Accordingly, this section of the SOC offers specific clinical guidelines for the assessment and treatment of gender dysphoric children and adolescents.

Differences between Children and Adolescents with Gender Dysphoria

An important difference between gender dysphoric children and adolescents is in the proportion for whom dysphoria persists into adulthood. Gender dysphoria during childhood does not inevitably continue into adulthood.⁵ Rather, in follow-up studies of prepubertal children (mainly boys) who were referred to clinics for assessment of gender dysphoria, the dysphoria persisted into adulthood for only 6-23% of children (Cohen-Kettenis, 2001; Zucker & Bradley, 1995). Boys in these studies were more likely to identify as gay in adulthood than as transgender (Green, 1987; Money & Russo, 1979; Zucker & Bradley, 1995; Zuger, 1984). Newer studies, also including girls, showed a 12-27% persistence rate of gender dysphoria into adulthood (Drummond, Bradley, Peterson-Badali, & Zucker, 2008; Wallien & Cohen-Kettenis, 2008).

In contrast, the persistence of gender dysphoria into adulthood appears to be much higher for adolescents. No formal prospective studies exist. However, in a follow-up study of 70 adolescents who were diagnosed with gender dysphoria and given puberty suppressing hormones, all continued with the actual sex reassignment, beginning with feminizing/masculinizing hormone therapy (de Vries, Steensma, Doreleijers, & Cohen-Kettenis, 2010).

Another difference between gender dysphoric children and adolescents is in the sex ratios for each age group. In clinically referred, gender dysphoric children under age 12, the male/female ratio ranges from 6:1 to 3:1 (Zucker, 2004). In clinically referred, gender dysphoric adolescents older than age 12, the male/female ratio is close to 1:1 (Cohen-Kettenis & Pfäfflin, 2003).

As discussed in section IV and by Zucker and Lawrence (2009), formal epidemiologic studies on gender dysphoria – in children, adolescents, and adults – are lacking. Additional research is needed to refine estimates of its prevalence and persistence in different populations worldwide.

⁵ Gender nonconforming behaviors in children may continue into adulthood, but such behaviors are not necessarily indicative of gender dysphoria and a need for treatment. As described in section III, gender dysphoria is not synonymous with diversity in gender expression.

Phenomenology in Children

Children as young as age two may show features that could indicate gender dysphoria. They may express a wish to be of the other sex and be unhappy about their physical sex characteristics and functions. In addition, they may prefer clothes, toys, and games that are commonly associated with the other sex and prefer playing with other-sex peers. There appears to be heterogeneity in these features: Some children demonstrate extremely gender nonconforming behavior and wishes, accompanied by persistent and severe discomfort with their primary sex characteristics. In other children, these characteristics are less intense or only partially present (Cohen-Kettenis et al., 2006; Knudson, De Cuypere, & Bockting, 2010a).

It is relatively common for gender dysphoric children to have co-existing internalizing disorders such as anxiety and depression (Cohen-Kettenis, Owen, Kaijser, Bradley, & Zucker, 2003; Wallien, Swaab, & Cohen-Kettenis, 2007; Zucker, Owen, Bradley, & Ameeriar, 2002). The prevalence of autistic spectrum disorders seems to be higher in clinically referred, gender dysphoric children than in the general population (de Vries, Noens, Cohen-Kettenis, van Berckelaer-Onnes, & Doreleijers, 2010).

Phenomenology in Adolescents

In most children, gender dysphoria will disappear before or early in puberty. However, in some children these feelings will intensify and body aversion will develop or increase as they become adolescents and their secondary sex characteristics develop (Cohen-Kettenis, 2001; Cohen-Kettenis & Pfäfflin, 2003; Drummond et al., 2008; Wallien & Cohen-Kettenis, 2008; Zucker & Bradley, 1995). Data from one study suggest that more extreme gender nonconformity in childhood is associated with persistence of gender dysphoria into late adolescence and early adulthood (Wallien & Cohen-Kettenis, 2008). Yet many adolescents and adults presenting with gender dysphoria do not report a history of childhood gender nonconforming behaviors (Docter, 1988; Landén, Wålinder, & Lundström, 1998). Therefore, it may come as a surprise to others (parents, other family members, friends, and community members) when a youth's gender dysphoria first becomes evident in adolescence.

Adolescents who experience their primary and/or secondary sex characteristics and their sex assigned at birth as inconsistent with their gender identity may be intensely distressed about it. Many, but not all, gender dysphoric adolescents have a strong wish for hormones and surgery. Increasing numbers of adolescents have already started living in their desired gender role upon entering high school (Cohen-Kettenis & Pfäfflin, 2003).

Among adolescents who are referred to gender identity clinics, the number considered eligible for early medical treatment – starting with GnRH analogues to suppress puberty in the first Tanner stages – differs among countries and centers. Not all clinics offer puberty suppression. If such treatment is offered, the pubertal stage at which adolescents are allowed to start varies from Tanner stage 2 to stage 4 (Delemarre-van de Waal & Cohen-Kettenis, 2006; Zucker et al., in press). The percentages of treated adolescents are likely influenced by the organization of health care, insurance aspects, cultural differences, opinions of health professionals, and diagnostic procedures offered in different settings.

Inexperienced clinicians may mistake indications of gender dysphoria for delusions. Phenomenologically, there is a qualitative difference between the presentation of gender dysphoria and the presentation of delusions or other psychotic symptoms. The vast majority of children and adolescents with gender dysphoria are not suffering from underlying severe psychiatric illness such as psychotic disorders (Steensma, Biemond, de Boer, & Cohen-Kettenis, published online ahead of print January 7, 2011).

It is more common for adolescents with gender dysphoria to have co-existing internalizing disorders such as anxiety and depression, and/or externalizing disorders such as oppositional defiant disorder (de Vries et al., 2010). As in children, there seems to be a higher prevalence of autistic spectrum disorders in clinically referred, gender dysphoric adolescents than in the general adolescent population (de Vries et al., 2010).

Competency of Mental Health Professionals Working with Children or Adolescents with Gender Dysphoria

The following are recommended minimum credentials for mental health professionals who assess, refer, and offer therapy to children and adolescents presenting with gender dysphoria:

1. Meet the competency requirements for mental health professionals working with adults, as outlined in section VII;
2. Trained in childhood and adolescent developmental psychopathology;
3. Competent in diagnosing and treating the ordinary problems of children and adolescents.

Roles of Mental Health Professionals Working with Children and Adolescents with Gender Dysphoria

The roles of mental health professionals working with gender dysphoric children and adolescents may include the following:

1. Directly assess gender dysphoria in children and adolescents (see general guidelines for assessment, below).
2. Provide family counseling and supportive psychotherapy to assist children and adolescents with exploring their gender identity, alleviating distress related to their gender dysphoria, and ameliorating any other psychosocial difficulties.
3. Assess and treat any co-existing mental health concerns of children or adolescents (or refer to another mental health professional for treatment). Such concerns should be addressed as part of the overall treatment plan.
4. Refer adolescents for additional physical interventions (such as puberty suppressing hormones) to alleviate gender dysphoria. The referral should include documentation of an assessment of gender dysphoria and mental health, the adolescent's eligibility for physical interventions (outlined below), the mental health professional's relevant expertise, and any other information pertinent to the youth's health and referral for specific treatments.
5. Educate and advocate on behalf of gender dysphoric children, adolescents, and their families in their community (e.g., day care centers, schools, camps, other organizations). This is particularly important in light of evidence that children and adolescents who do not conform to socially prescribed gender norms may experience harassment in school (Grossman, D'Augelli, & Salter, 2006; Grossman, D'Augelli, Howell, & Hubbard, 2006; Sausa, 2005), putting them at risk for social isolation, depression, and other negative sequelae (Nuttbrock et al., 2010).
6. Provide children, youth, and their families with information and referral for peer support, such as support groups for parents of gender nonconforming and transgender children (Gold & MacNish, 2011; Pleak, 1999; Rosenberg, 2002).

Assessment and psychosocial interventions for children and adolescents are often provided within a multi-disciplinary gender identity specialty service. If such a multidisciplinary service is not available, a mental health professional should provide consultation and liaison arrangements with a pediatric endocrinologist for the purpose of assessment, education, and involvement in any decisions about physical interventions.

Psychological Assessment of Children and Adolescents

When assessing children and adolescents who present with gender dysphoria, mental health professionals should broadly conform to the following guidelines:

1. Mental health professionals should not dismiss or express a negative attitude towards nonconforming gender identities or indications of gender dysphoria. Rather, they should acknowledge the presenting concerns of children, adolescents, and their families; offer a thorough assessment for gender dysphoria and any co-existing mental health concerns; and educate clients and their families about therapeutic options, if needed. Acceptance and removal of secrecy can bring considerable relief to gender dysphoric children/adolescents and their families.
2. Assessment of gender dysphoria and mental health should explore the nature and characteristics of a child's or adolescent's gender identity. A psychodiagnostic and psychiatric assessment – covering the areas of emotional functioning, peer and other social relationships, and intellectual functioning/school achievement – should be performed. Assessment should include an evaluation of the strengths and weaknesses of family functioning. Emotional and behavioral problems are relatively common, and unresolved issues in a child's or youth's environment may be present (de Vries, Doreleijers, Steensma, & Cohen-Kettenis, 2011; Di Ceglie & Thümmel, 2006; Wallien et al., 2007).
3. For adolescents, the assessment phase should also be used to inform youth and their families about the possibilities and limitations of different treatments. This is necessary for informed consent, but also important for assessment. The way that adolescents respond to information about the reality of sex reassignment can be diagnostically informative. Correct information may alter a youth's desire for certain treatment, if the desire was based on unrealistic expectations of its possibilities.

Psychological and Social Interventions for Children and Adolescents

When supporting and treating children and adolescents with gender dysphoria, health professionals should broadly conform to the following guidelines:

1. Mental health professionals should help families to have an accepting and nurturing response to the concerns of their gender dysphoric child or adolescent. Families play an important role in the psychological health and well-being of youth (Brill & Pepper, 2008; Lev, 2004). This also applies to peers and mentors from the community, who can be another source of social support.

2. Psychotherapy should focus on reducing a child's or adolescent's distress related to the gender dysphoria and on ameliorating any other psychosocial difficulties. For youth pursuing sex reassignment, psychotherapy may focus on supporting them before, during, and after reassignment. Formal evaluations of different psychotherapeutic approaches for this situation have not been published, but several counseling methods have been described (Cohen-Kettenis, 2006; de Vries, Cohen-Kettenis, & Delemarre-van de Waal, 2006; Di Ceglie & Thümmel, 2006; Hill, Menvielle, Sica, & Johnson, 2010; Malpas, in press; Menvielle & Tuerk, 2002; Rosenberg, 2002; Vanderburgh, 2009; Zucker, 2006).

Treatment aimed at trying to change a person's gender identity and expression to become more congruent with sex assigned at birth has been attempted in the past without success (Gelder & Marks, 1969; Greenson, 1964), particularly in the long term (Cohen-Kettenis & Kuiper, 1984; Pauly, 1965). Such treatment is no longer considered ethical.

1. Families should be supported in managing uncertainty and anxiety about their child's or adolescent's psychosexual outcomes and in helping youth to develop a positive self-concept.
2. Mental health professionals should not impose a binary view of gender. They should give ample room for clients to explore different options for gender expression. Hormonal or surgical interventions are appropriate for some adolescents, but not for others.
3. Clients and their families should be supported in making difficult decisions regarding the extent to which clients are allowed to express a gender role that is consistent with their gender identity, as well as the timing of changes in gender role and possible social transition. For example, a client might attend school while undergoing social transition only partly (e.g., by wearing clothing and having a hairstyle that reflects gender identity) or completely (e.g., by also using a name and pronouns congruent with gender identity). Difficult issues include whether and when to inform other people of the client's situation, and how others in their lives should respond.
4. Health professionals should support clients and their families as educators and advocates in their interactions with community members and authorities such as teachers, school boards, and courts.
5. Mental health professionals should strive to maintain a therapeutic relationship with gender nonconforming children/adolescents and their families throughout any subsequent social changes or physical interventions. This ensures that decisions about gender expression and the treatment of gender dysphoria are thoughtfully and recurrently considered. The same reasoning applies if a child or adolescent has already socially changed gender role prior to being seen by a mental health professional.

Social Transition in Early Childhood

Some children state that they want to make a social transition to a different gender role long before puberty. For some children, this may reflect an expression of their gender identity. For others, this could be motivated by other forces. Families vary in the extent to which they allow their young children to make a social transition to another gender role. Social transitions in early childhood do occur within some families with early success. This is a controversial issue, and divergent views are held by health professionals. The current evidence base is insufficient to predict the long-term outcomes of completing a gender role transition during early childhood. Outcomes research with children who completed early social transitions would greatly inform future clinical recommendations.

Mental health professionals can help families to make decisions regarding the timing and process of any gender role changes for their young children. They should provide information and help parents to weigh the potential benefits and challenges of particular choices. Relevant in this respect are the previously described relatively low persistence rates of childhood gender dysphoria (Drummond et al., 2008; Wallien & Cohen-Kettenis, 2008). A change back to the original gender role can be highly distressing and even result in postponement of this second social transition on the child's part (Steensma & Cohen-Kettenis, 2011). For reasons such as these, parents may want to present this role change as an exploration of living in another gender role, rather than an irreversible situation. Mental health professionals can assist parents in identifying potential in-between solutions or compromises (e.g., only when on vacation). It is also important that parents explicitly let the child know that there is a way back.

Regardless of a family's decisions regarding transition (timing, extent), professionals should counsel and support them as they work through the options and implications. If parents do not allow their young child to make a gender role transition, they may need counseling to assist them with meeting their child's needs in a sensitive and nurturing way, ensuring that the child has ample possibilities to explore gender feelings and behavior in a safe environment. If parents do allow their young child to make a gender role transition, they may need counseling to facilitate a positive experience for their child. For example, they may need support in using correct pronouns, maintaining a safe and supportive environment for their transitioning child (e.g., in school, peer group settings), and communicating with other people in their child's life. In either case, as a child nears puberty, further assessment may be needed as options for physical interventions become relevant.

Physical Interventions for Adolescents

Before any physical interventions are considered for adolescents, extensive exploration of psychological, family, and social issues should be undertaken, as outlined above. The duration of this exploration may vary considerably depending on the complexity of the situation.

Physical interventions should be addressed in the context of adolescent development. Some identity beliefs in adolescents may become firmly held and strongly expressed, giving a false impression of irreversibility. An adolescent's shift towards gender conformity can occur primarily to please the parents and may not persist or reflect a permanent change in gender dysphoria (Hembree et al., 2009; Steensma et al., published online ahead of print January 7, 2011).

Physical interventions for adolescents fall into three categories or stages (Hembree et al., 2009):

1. *Fully reversible interventions.* These involve the use of GnRH analogues to suppress estrogen or testosterone production and consequently delay the physical changes of puberty. Alternative treatment options include progestins (most commonly medroxyprogesterone) or other medications (such as spironolactone) that decrease the effects of androgens secreted by the testicles of adolescents who are not receiving GnRH analogues. Continuous oral contraceptives (or depot medroxyprogesterone) may be used to suppress menses.
2. *Partially reversible interventions.* These include hormone therapy to masculinize or feminize the body. Some hormone-induced changes may need reconstructive surgery to reverse the effect (e.g., gynaecomastia caused by estrogens), while other changes are not reversible (e.g., deepening of the voice caused by testosterone).
3. *Irreversible interventions.* These are surgical procedures.

A staged process is recommended to keep options open through the first two stages. Moving from one stage to another should not occur until there has been adequate time for adolescents and their parents to assimilate fully the effects of earlier interventions.

Fully Reversible Interventions

Adolescents may be eligible for puberty suppressing hormones as soon as pubertal changes have begun. In order for adolescents and their parents to make an informed decision about pubertal delay, it is recommended that adolescents experience the onset of puberty to at least Tanner Stage 2. Some children may arrive at this stage at very young ages (e.g., 9 years of age). Studies

evaluating this approach only included children who were at least 12 years of age (Cohen-Kettenis, Schagen, Steensma, de Vries, & Delemarre-van de Waal, 2011; de Vries, Steensma et al., 2010; Delemarre-van de Waal, van Weissenbruch, & Cohen Kettenis, 2004; Delemarre-van de Waal & Cohen-Kettenis, 2006).

Two goals justify intervention with puberty suppressing hormones: (i) their use gives adolescents more time to explore their gender nonconformity and other developmental issues; and (ii) their use may facilitate transition by preventing the development of sex characteristics that are difficult or impossible to reverse if adolescents continue on to pursue sex reassignment.

Puberty suppression may continue for a few years, at which time a decision is made to either discontinue all hormone therapy or transition to a feminizing/masculinizing hormone regimen. Pubertal suppression does not inevitably lead to social transition or to sex reassignment.

Criteria for puberty suppressing hormones

In order for adolescents to receive puberty suppressing hormones, the following minimum criteria must be met:

1. The adolescent has demonstrated a long-lasting and intense pattern of gender nonconformity or gender dysphoria (whether suppressed or expressed);
2. Gender dysphoria emerged or worsened with the onset of puberty;
3. Any co-existing psychological, medical, or social problems that could interfere with treatment (e.g., that may compromise treatment adherence) have been addressed, such that the adolescent's situation and functioning are stable enough to start treatment;
4. The adolescent has given informed consent and, particularly when the adolescent has not reached the age of medical consent, the parents or other caretakers or guardians have consented to the treatment and are involved in supporting the adolescent throughout the treatment process.

Regimens, monitoring, and risks for puberty suppression

For puberty suppression, adolescents with male genitalia should be treated with GnRH analogues, which stop luteinizing hormone secretion and therefore testosterone secretion. Alternatively, they may be treated with progestins (such as medroxyprogesterone) or with other medications that block testosterone secretion and/or neutralize testosterone action. Adolescents with female genitalia should be treated with GnRH analogues, which stop the production of estrogens and

progesterone. Alternatively, they may be treated with progestins (such as medroxyprogesterone). Continuous oral contraceptives (or depot medroxyprogesterone) may be used to suppress menses. In both groups of adolescents, use of GnRH analogues is the preferred treatment (Hembree et al., 2009), but their high cost is prohibitive for some patients

During pubertal suppression, an adolescent's physical development should be carefully monitored – preferably by a pediatric endocrinologist – so that any necessary interventions can occur (e.g., to establish an adequate gender appropriate height, to improve iatrogenic low bone marrow density) (Hembree et al., 2009).

Early use of puberty suppressing hormones may avert negative social and emotional consequences of gender dysphoria more effectively than their later use would. Intervention in early adolescence should be managed with pediatric endocrinological advice, when available. Adolescents with male genitalia who start GnRH analogues early in puberty should be informed that this could result in insufficient penile tissue for penile inversion vaginoplasty techniques (alternative techniques, such as the use of a skin graft or colon tissue, are available).

Neither puberty suppression nor allowing puberty to occur is a neutral act. On the one hand, functioning in later life can be compromised by the development of irreversible secondary sex characteristics during puberty and by years spent experiencing intense gender dysphoria. On the other hand, there are concerns about negative physical side effects of GnRH analog use (e.g., on bone development and height). Although the very first results of this approach (as assessed for adolescents followed over 10 years) are promising (Cohen-Kettenis et al., 2011; Delemarre-van de Waal & Cohen-Kettenis, 2006), the long-term effects can only be determined when the earliest treated patients reach the appropriate age.

Partially Reversible Interventions

Adolescents may be eligible to begin feminizing/masculinizing hormone therapy, preferably with parental consent. In many countries, 16-year-olds are legal adults for medical decision-making and do not require parental consent. Ideally, treatment decisions should be made among the adolescent, the family, and the treatment team.

Regimens for hormone therapy in gender dysphoric adolescents differ substantially from those used in adults (Hembree et al., 2009). The hormone regimens for youth are adapted to account for the somatic, emotional, and mental development that occurs throughout adolescence (Hembree et al., 2009).

Irreversible Interventions

Genital surgery should not be carried out until (i) patients reach the legal age of majority in a given country, and (ii) patients have lived continuously for at least 12 months in the gender role that is congruent with their gender identity. The age threshold should be seen as a minimum criterion and not an indication in and of itself for active intervention.

Chest surgery in FtM patients could be carried out earlier, preferably after ample time of living in the desired gender role and after one year of testosterone treatment. The intent of this suggested sequence is to give adolescents sufficient opportunity to experience and socially adjust in a more masculine gender role, before undergoing irreversible surgery. However, different approaches may be more suitable, depending on an adolescent's specific clinical situation and goals for gender identity expression.

Risks of Withholding Medical Treatment for Adolescents

Refusing timely medical interventions for adolescents might prolong gender dysphoria and contribute to an appearance that could provoke abuse and stigmatization. As the level of gender-related abuse is strongly associated with the degree of psychiatric distress during adolescence (Nuttbrock et al., 2010), withholding puberty suppression and subsequent feminizing or masculinizing hormone therapy is not a neutral option for adolescents.

VII

Mental Health

Transsexual, transgender, and gender nonconforming people might seek the assistance of a mental health professional for any number of reasons. Regardless of a person's reason for seeking care, mental health professionals should have familiarity with gender nonconformity, act with appropriate cultural competence, and exhibit sensitivity in providing care.

This section of the *SOC* focuses on the role of mental health professionals in the care of adults seeking help for gender dysphoria and related concerns. Professionals working with gender dysphoric children, adolescents, and their families should consult section VI.

Issues of Access to Care

Qualified mental health professionals are not universally available; thus, access to quality care might be limited. WPATH aims to improve access and provides regular continuing education opportunities to train professionals from various disciplines to provide quality, transgender-specific health care. Providing mental health care from a distance through the use of technology may be one way to improve access (Fraser, 2009b).

In many places around the world, access to health care for transsexual, transgender, and gender nonconforming people is also limited by a lack of health insurance or other means to pay for needed care. WPATH urges health insurance companies and other third-party payers to cover the medically necessary treatment to alleviate gender dysphoria (American Medical Association, 2008; Anton, 2009; The World Professional Association for Transgender Health, 2008).

When faced with a client who is unable to access services, referral to available peer support resources (offline and online) is recommended. Finally, harm reduction approaches might be indicated to assist clients with making healthy decisions to improve their lives.

VIII

Hormone Therapy

Medical Necessity of Hormone Therapy

Feminizing/masculinizing hormone therapy – the administration of exogenous endocrine agents to induce feminizing or masculinizing changes – is a medically necessary intervention for many transsexual, transgender, and gender nonconforming individuals with gender dysphoria (Newfield, Hart, Dibble, & Kohler, 2006; Pfäfflin & Junge, 1998). Some people seek maximum feminization/masculinization, while others experience relief with an androgynous presentation resulting from hormonal minimization of existing secondary sex characteristics (Factor & Rothblum, 2008). Evidence for the psychosocial outcomes of hormone therapy is summarized in Appendix D.

Hormone therapy must be individualized based on a patient's goals, the risk/benefit ratio of medications, the presence of other medical conditions, and consideration of social and economic issues. Hormone therapy can provide significant comfort to patients who do not wish to make a social gender role transition or undergo surgery, or who are unable to do so (Meyer III, 2009).

Hormone therapy is a recommended criterion for some, but not all, surgical treatments for gender dysphoria (see section XI and Appendix C).

Criteria for Hormone Therapy

Initiation of hormone therapy may be undertaken after a psychosocial assessment has been conducted and informed consent has been obtained by a qualified health professional, as outlined in section VII of the SOC. A referral is required from the mental health professional who performed the assessment, unless the assessment was done by a hormone provider who is also qualified in this area.

The criteria for hormone therapy are as follows:

1. Persistent, well-documented gender dysphoria;
2. Capacity to make a fully informed decision and to consent for treatment;
3. Age of majority in a given country (if younger, follow the *Standards of Care* outlined in section VI);
4. If significant medical or mental health concerns are present, they must be reasonably well-controlled.

As noted in section VII of the SOC, the presence of co-existing mental health concerns does not necessarily preclude access to feminizing/masculinizing hormones; rather, these concerns need to be managed prior to or concurrent with treatment of gender dysphoria.

In selected circumstances, it can be acceptable practice to provide hormones to patients who have not fulfilled these criteria. Examples include facilitating the provision of monitored therapy using hormones of known quality as an alternative to illicit or unsupervised hormone use or to patients who have already established themselves in their affirmed gender and who have a history of prior hormone use. It is unethical to deny availability or eligibility for hormone therapy solely on the basis of blood seropositivity for blood-borne infections such as HIV or hepatitis B or C.

In rare cases, hormone therapy may be contraindicated due to serious individual health conditions. Health professionals should assist these patients with accessing non-hormonal interventions for gender dysphoria. A qualified mental health professional familiar with the patient is an excellent resource in these circumstances.

Informed Consent

Feminizing/masculinizing hormone therapy may lead to irreversible physical changes. Thus, hormone therapy should be provided only to those who are legally able to provide informed consent. This includes people who have been declared by a court to be emancipated minors, incarcerated people, and cognitively impaired people who are considered competent to participate in their medical decisions (see also Bockting et al., 2006). Providers should document in the medical record that comprehensive information has been provided and understood about all relevant aspects of the hormone therapy, including both possible benefits and risks and the impact on reproductive capacity.

Relationship between the Standards of Care and Informed Consent Model Protocols

A number of community health centers in the United States have developed protocols for providing hormone therapy based on an approach that has become known as the Informed Consent Model (Callen Lorde Community Health Center, 2000, 2011; Fenway Community Health Transgender Health Program, 2007; Tom Waddell Health Center, 2006). These protocols are consistent with the guidelines presented in the WPATH *Standards of Care, Version 7*. The SOC are flexible clinical guidelines; they allow for tailoring of interventions to the needs of the individual receiving services and for tailoring of protocols to the approach and setting in which these services are provided (Ehrbar & Gorton, 2010).

Obtaining informed consent for hormone therapy is an important task of providers to ensure that patients understand the psychological and physical benefits and risks of hormone therapy, as well as its psychosocial implications. Providers prescribing the hormones or health professionals recommending the hormones should have the knowledge and experience to assess gender dysphoria. They should inform individuals of the particular benefits, limitations, and risks of hormones, given the patient's age, previous experience with hormones, and concurrent physical or mental health concerns.

Screening for and addressing acute or current mental health concerns is an important part of the informed consent process. This may be done by a mental health professional or by an appropriately trained prescribing provider (see section VII of the SOC). The same provider or another appropriately trained member of the health care team (e.g., a nurse) can address the psychosocial implications of taking hormones when necessary (e.g., the impact of masculinization/feminization on how one is perceived and its potential impact on relationships with family, friends, and coworkers). If indicated, these providers will make referrals for psychotherapy and for the assessment and treatment of co-existing mental health concerns such as anxiety or depression.

The difference between the Informed Consent Model and *SOC, Version 7* is that the *SOC* puts greater emphasis on the important role that mental health professionals can play in alleviating gender dysphoria and facilitating changes in gender role and psychosocial adjustment. This may include a comprehensive mental health assessment and psychotherapy, when indicated. In the Informed Consent Model, the focus is on obtaining informed consent as the threshold for the initiation of hormone therapy in a multidisciplinary, harm-reduction environment. Less emphasis is placed on the provision of mental health care until the patient requests it, unless significant mental health concerns are identified that would need to be addressed before hormone prescription.

Physical Effects of Hormone Therapy

Feminizing/masculinizing hormone therapy will induce physical changes that are more congruent with a patient's gender identity.

- In FtM patients, the following physical changes are expected to occur: deepened voice, clitoral enlargement (variable), growth in facial and body hair, cessation of menses, atrophy of breast tissue, increased libido, and decreased percentage of body fat compared to muscle mass.
- In MtF patients, the following physical changes are expected to occur: breast growth (variable), decreased libido and erections, decreased testicular size, and increased percentage of body fat compared to muscle mass.

Most physical changes, whether feminizing or masculinizing, occur over the course of two years. The amount of physical change and the exact timeline of effects can be highly variable. Tables 1a and 1b outline the approximate time course of these physical changes.

TABLE 1A: EFFECTS AND EXPECTED TIME COURSE OF MASCULINIZING HORMONES^A

Effect	Expected Onset ^B	Expected Maximum Effect ^B
Skin oiliness/acne	1-6 months	1-2 years
Facial/body hair growth	3-6 months	3-5 years
Scalp hair loss	>12 months ^C	variable
Increased muscle mass/strength	6-12 months	2-5 years ^D
Body fat redistribution	3-6 months	2-5 years
Cessation of menses	2-6 months	n/a
Clitoral enlargement	3-6 months	1-2 years
Vaginal atrophy	3-6 months	1-2 years
Deepened voice	3-12 months	1-2 years

^A Adapted with permission from Hembree et al.(2009). Copyright 2009, The Endocrine Society.

^B Estimates represent published and unpublished clinical observations.

^C Highly dependent on age and inheritance; may be minimal.

^D Significantly dependent on amount of exercise.

TABLE 1B: EFFECTS AND EXPECTED TIME COURSE OF FEMINIZING HORMONES^A

Effect	Expected Onset ^B	Expected Maximum Effect ^B
Body fat redistribution	3-6 months	2-5 years
Decreased muscle mass/ strength	3-6 months	1-2 years ^C
Softening of skin/decreased oiliness	3-6 months	unknown
Decreased libido	1-3 months	1-2 years
Decreased spontaneous erections	1-3 months	3-6 months
Male sexual dysfunction	variable	variable
Breast growth	3-6 months	2-3 years
Decreased testicular volume	3-6 months	2-3 years
Decreased sperm production	variable	variable
Thinning and slowed growth of body and facial hair	6-12 months	> 3 years ^D
Male pattern baldness	No regrowth, loss stops 1-3 months	1-2 years

^A Adapted with permission from Hembree et al. (2009). Copyright 2009, *The Endocrine Society*.

^B Estimates represent published and unpublished clinical observations.

^C Significantly dependent on amount of exercise.

^D Complete removal of male facial and body hair requires electrolysis, laser treatment, or both.

The degree and rate of physical effects depends in part on the dose, route of administration, and medications used, which are selected in accordance with a patient's specific medical goals (e.g., changes in gender role expression, plans for sex reassignment) and medical risk profile. There is no current evidence that response to hormone therapy – with the possible exception of voice deepening in FtM persons – can be reliably predicted based on age, body habitus, ethnicity, or family appearance. All other factors being equal, there is no evidence to suggest that any medically approved type or method of administering hormones is more effective than any other in producing the desired physical changes.

Risks of Hormone Therapy

All medical interventions carry risks. The likelihood of a serious adverse event is dependent on numerous factors: the medication itself, dose, route of administration, and a patient's clinical characteristics (age, co-morbidities, family history, health habits). It is thus impossible to predict whether a given adverse effect will happen in an individual patient.

The risks associated with feminizing/masculinizing hormone therapy for the transsexual, transgender, and gender nonconforming population as a whole are summarized in Table 2. Based on the level of evidence, risks are categorized as follows: (i) likely increased risk with hormone therapy, (ii) possibly increased risk with hormone therapy, or (iii) inconclusive or no increased risk. Items in the last category include those that may present risk, but for which the evidence is so minimal that no clear conclusion can be reached.

Additional detail about these risks can be found in Appendix B, which is based on two comprehensive, evidence-based literature reviews of masculinizing/feminizing hormone therapy (Feldman & Safer, 2009; Hembree et al., 2009), along with a large cohort study (Asscheman et al., 2011). These reviews can serve as detailed references for providers, along with other widely recognized, published clinical materials (Dahl, Feldman, Goldberg, & Jaber, 2006; Ettner, Monstrey, & Eyler, 2007).

TABLE 2: RISKS ASSOCIATED WITH HORMONE THERAPY. BOLDDED ITEMS ARE CLINICALLY SIGNIFICANT

Risk Level	Feminizing hormones	Masculinizing hormones
Likely increased risk	Venous thromboembolic disease^A Gallstones Elevated liver enzymes Weight gain Hypertriglyceridemia	Polycythemia Weight gain Acne Androgenic alopecia (balding) Sleep apnea
Likely increased risk with presence of additional risk factors ^B	Cardiovascular disease	
Possible increased risk	Hypertension Hyperprolactinemia or prolactinoma ^A	Elevated liver enzymes Hyperlipidemia
Possible increased risk with presence of additional risk factors ^B	Type 2 diabetes^A	Destabilization of certain psychiatric disorders^C Cardiovascular disease Hypertension Type 2 diabetes
No increased risk or inconclusive	Breast cancer	Loss of bone density Breast cancer Cervical cancer Ovarian cancer Uterine cancer

^A Risk is greater with oral estrogen administration than with transdermal estrogen administration.

^B Additional risk factors include age.

^C Includes bipolar, schizoaffective, and other disorders that may include manic or psychotic symptoms. This adverse event appears to be associated with higher doses or supraphysiologic blood levels of testosterone.

Competency of Hormone-Prescribing Physicians, Relationship with Other Health Professionals

Feminizing/masculinizing hormone therapy is best undertaken in the context of a complete approach to health care that includes comprehensive primary care and a coordinated approach to psychosocial issues (Feldman & Safer, 2009). While psychotherapy or ongoing counseling is not required for the initiation of hormone therapy, if a therapist is involved, then regular communication among health professionals is advised (with the patient's consent) to ensure that the transition process is going well, both physically and psychosocially.

With appropriate training, feminizing/masculinizing hormone therapy can be managed by a variety of providers, including nurse practitioners and primary care physicians (Dahl et al., 2006). Medical visits relating to hormone maintenance provide an opportunity to deliver broader care to a population that is often medically underserved (Clements, Wilkinson, Kitano, & Marx, 1999; Feldman, 2007; Xavier, 2000). Many of the screening tasks and management of co-morbidities associated with long-term hormone use, such as cardiovascular risk factors and cancer screening, fall more uniformly within the scope of primary care rather than specialist care (American Academy of Family Physicians, 2005; Eyer, 2007; World Health Organization, 2008), particularly in locations where dedicated gender teams or specialized physicians are not available.

Given the multidisciplinary needs of transsexual, transgender, and gender nonconforming people seeking hormone therapy, as well as the difficulties associated with fragmentation of care in general (World Health Organization, 2008), WPATH strongly encourages the increased training and involvement of primary care providers in the area of feminizing/masculinizing hormone therapy. If hormones are prescribed by a specialist, there should be close communication with the patient's primary care provider. Conversely, an experienced hormone provider or endocrinologist should be involved if the primary care physician has no experience with this type of hormone therapy, or if the patient has a pre-existing metabolic or endocrine disorder that could be affected by endocrine therapy.

While formal training programs in transgender medicine do not yet exist, hormone providers have a responsibility to obtain appropriate knowledge and experience in this field. Clinicians can increase their experience and comfort in providing feminizing/masculinizing hormone therapy by co-managing care or consulting with a more experienced provider, or by providing more limited types of hormone therapy before progressing to initiation of hormone therapy. Because this field of medicine is evolving, clinicians should become familiar and keep current with the medical literature, and discuss emerging issues with colleagues. Such discussions might occur through networks established by WPATH and other national/local organizations.

Responsibilities of Hormone-Prescribing Physicians

In general, clinicians who prescribe hormone therapy should engage in the following tasks:

1. Perform an initial evaluation that includes discussion of a patient's physical transition goals, health history, physical examination, risk assessment, and relevant laboratory tests.
2. Discuss with patients the expected effects of feminizing/masculinizing medications and the possible adverse health effects. These effects can include a reduction in fertility (Feldman & Safer, 2009; Hembree et al., 2009). Therefore, reproductive options should be discussed with patients before starting hormone therapy (see section IX).
3. Confirm that patients have the capacity to understand the risks and benefits of treatment and are capable of making an informed decision about medical care.
4. Provide ongoing medical monitoring, including regular physical and laboratory examination to monitor hormone effectiveness and side effects.
5. Communicate as needed with a patient's primary care provider, mental health professional, and surgeon.
6. If needed, provide patients with a brief written statement indicating that they are under medical supervision and care that includes feminizing/masculinizing hormone therapy. Particularly during the early phases of hormone treatment, a patient may wish to carry this statement at all times to help prevent difficulties with the police and other authorities.

Depending on the clinical situation for providing hormones (see below), some of these responsibilities are less relevant. Thus, the degree of counseling, physical examinations, and laboratory evaluations should be individualized to a patient's needs.

Clinical Situations for Hormone Therapy

There are circumstances in which clinicians may be called upon to provide hormones without necessarily initiating or maintaining long-term feminizing/masculinizing hormone therapy. By acknowledging these different clinical situations (see below, from least to highest level of complexity), it may be possible to involve clinicians in feminizing/masculinizing hormone therapy who might not otherwise feel able to offer this treatment.

1. Bridging

Whether prescribed by another clinician or obtained through other means (e.g., purchased over the internet), patients may present for care already on hormone therapy. Clinicians can provide a limited (1-6 month) prescription for hormones while helping patients find a provider who can prescribe long-term hormone therapy. Providers should assess a patient's current regimen for safety and drug interactions and substitute safer medications or doses when indicated (Dahl et al., 2006; Feldman & Safer, 2009). If hormones were previously prescribed, medical records should be requested (with the patient's permission) to obtain the results of baseline examinations and laboratory tests and any adverse events. Hormone providers should also communicate with any mental health professional who is currently involved in a patient's care. If a patient has never had a psychosocial assessment as recommended by the SOC (see section VII), clinicians should refer the patient to a qualified mental health professional if appropriate and feasible (Feldman & Safer, 2009). Providers who prescribe bridging hormones need to work with patients to establish limits as to the duration of bridging therapy.

2. Hormone therapy following gonad removal

Hormone replacement with estrogen or testosterone is usually continued lifelong after an oophorectomy or orchiectomy, unless medical contraindications arise. Because hormone doses are often decreased after these surgeries (Basson, 2001; Levy, Crown, & Reid, 2003; Moore, Wisniewski, & Dobs, 2003) and only adjusted for age and co-morbid health concerns, hormone management in this situation is quite similar to hormone replacement in any hypogonadal patient.

3. Hormone maintenance prior to gonad removal

Once patients have achieved maximal feminizing/masculinizing benefits from hormones (typically two or more years), they remain on a maintenance dose. The maintenance dose is then adjusted for changes in health conditions, aging, or other considerations such as lifestyle changes (Dahl et al., 2006). When a patient on maintenance hormones presents for care, the provider should assess the patient's current regimen for safety and drug interactions and substitute safer medications or doses when indicated. The patient should continue to be monitored by physical examinations and laboratory testing on a regular basis, as outlined in the literature (Feldman & Safer, 2009; Hembree et al., 2009). The dose and form of hormones should be revisited regularly with any changes in the patient's health status and available evidence on the potential long-term risks of hormones (See *Hormone Regimens*, below).

4. Initiating hormonal feminization/masculinization

This clinical situation requires the greatest commitment in terms of provider time and expertise. Hormone therapy must be individualized based on a patient's goals, the risk/benefit ratio of medications, the presence of other medical conditions, and consideration of social and economic issues. Although a wide variety of hormone regimens have been published (Dahl et al., 2006; Hembree et al., 2009; Moore et al., 2003), there are no published reports of randomized clinical trials comparing safety and efficacy. Despite this variation, a reasonable framework for initial risk assessment and ongoing monitoring of hormone therapy can be constructed, based on the efficacy and safety evidence presented above.

Risk Assessment and Modification for Initiating Hormone Therapy

The initial evaluation for hormone therapy assesses a patient's clinical goals and risk factors for hormone-related adverse events. During the risk assessment, the patient and clinician should develop a plan for reducing risks wherever possible, either prior to initiating therapy or as part of ongoing harm reduction.

All assessments should include a thorough physical exam, including weight, height, and blood pressure. The need for breast, genital, and rectal exams, which are sensitive issues for most transsexual, transgender, and gender nonconforming patients, should be based on individual risks and preventive health care needs (Feldman & Goldberg, 2006; Feldman, 2007).

Preventive care

Hormone providers should address preventive health care with patients, particularly if a patient does not have a primary care provider. Depending on a patient's age and risk profile, there may be appropriate screening tests or exams for conditions affected by hormone therapy. Ideally, these screening tests should be carried out prior to the start of hormone therapy.

Risk assessment and modification for feminizing hormone therapy (MtF)

There are no absolute contraindications to feminizing therapy *per se*, but absolute contraindications exist for the different feminizing agents, particularly estrogen. These include previous venous thrombotic events related to an underlying hypercoagulable condition, history of estrogen-sensitive neoplasm, and end-stage chronic liver disease (Gharib et al., 2005).

Other medical conditions, as noted in Table 2 and Appendix B, can be exacerbated by estrogen or androgen blockade, and therefore should be evaluated and reasonably well controlled prior to starting hormone therapy (Feldman & Safer, 2009; Hembree et al., 2009). Clinicians should particularly attend to tobacco use, as it is associated with increased risk of venous thrombosis, which is further increased with estrogen use. Consultation with a cardiologist may be advisable for patients with known cardio- or cerebrovascular disease.

Baseline laboratory values are important to both assess initial risk and evaluate possible future adverse events. Initial labs should be based on the risks of feminizing hormone therapy outlined in Table 2, as well as individual patient risk factors, including family history. Suggested initial lab panels have been published (Feldman & Safer, 2009; Hembree et al., 2009). These can be modified for patients or health care systems with limited resources, and in otherwise healthy patients.

Risk assessment and modification for masculinizing hormone therapy (FtM)

Absolute contraindications to testosterone therapy include pregnancy, unstable coronary artery disease, and untreated polycythemia with a hematocrit of 55% or higher (Carnegie, 2004). Because the aromatization of testosterone to estrogen may increase risk in patients with a history of breast or other estrogen dependent cancers (Moore et al., 2003), consultation with an oncologist may be indicated prior to hormone use. Co-morbid conditions likely to be exacerbated by testosterone use should be evaluated and treated, ideally prior to starting hormone therapy (Feldman & Safer, 2009; Hembree et al., 2009). Consultation with a cardiologist may be advisable for patients with known cardio- or cerebrovascular disease.

An increased prevalence of polycystic ovarian syndrome (PCOS) has been noted among FtM patients even in the absence of testosterone use (Baba et al., 2007; Balen, Schachter, Montgomery, Reid, & Jacobs, 1993; Bosinski et al., 1997). While there is no evidence that PCOS is related to the development of a transsexual, transgender, or gender nonconforming identity, PCOS is associated with increased risk of diabetes, cardiac disease, high blood pressure, and ovarian and endometrial cancers (Catrall & Healy, 2004). Signs and symptoms of PCOS should be evaluated prior to initiating testosterone therapy, as testosterone may affect many of these conditions. Testosterone can affect the developing fetus (Physicians' Desk Reference, 2011), and patients at risk of becoming pregnant require highly effective birth control.

Baseline laboratory values are important to both assess initial risk and evaluate possible future adverse events. Initial labs should be based on the risks of masculinizing hormone therapy outlined in Table 2, as well as individual patient risk factors, including family history. Suggested initial lab panels have been published (Feldman & Safer, 2009; Hembree et al., 2009). These can be modified for patients or health care systems with limited resources, and in otherwise healthy patients.

Clinical Monitoring during Hormone Therapy for Efficacy and Adverse Events

The purpose of clinical monitoring during hormone use is to assess the degree of feminization/masculinization and the possible presence of adverse effects of medication. However, as with the monitoring of any long-term medication, monitoring should take place in the context of comprehensive health care. Suggested clinical monitoring protocols have been published (Feldman & Safer, 2009; Hembree et al., 2009). Patients with co-morbid medical conditions may need to be monitored more frequently. Healthy patients in geographically remote or resource-poor areas may be able to use alternative strategies, such as telehealth, or cooperation with local providers such as nurses and physician assistants. In the absence of other indications, health professionals may prioritize monitoring for those risks that are either likely to be increased by hormone therapy or possibly increased by hormone therapy but clinically serious in nature.

Efficacy and risk monitoring during feminizing hormone therapy (MtF)

The best assessment of hormone efficacy is clinical response: Is a patient developing a feminized body while minimizing masculine characteristics, consistent with that patient's gender goals? In order to more rapidly predict the hormone dosages that will achieve clinical response, one can measure testosterone levels for suppression below the upper limit of the normal female range, and estradiol levels within a premenopausal female range but well below supraphysiologic levels (Feldman & Safer, 2009; Hembree et al., 2009).

Monitoring for adverse events should include both clinical and laboratory evaluation. Follow-up should include careful assessment for signs of cardiovascular impairment and venous thromboembolism (VTE) through measurement of blood pressure, weight, and pulse; heart and lung exams; and examination of the extremities for peripheral edema, localized swelling, or pain (Feldman & Safer, 2009). Laboratory monitoring should be based on the risks of hormone therapy described above, a patient's individual co-morbidities and risk factors, and the specific hormone regimen itself. Specific lab monitoring protocols have been published (Feldman & Safer, 2009; Hembree et al., 2009).

Efficacy and risk monitoring during masculinizing hormone therapy (FtM)

The best assessment of hormone efficacy is clinical response: Is a patient developing a masculinized body while minimizing feminine characteristics, consistent with that patient's gender goals? Clinicians can achieve a good clinical response with the least likelihood of adverse events by maintaining testosterone levels within the normal male range while avoiding supraphysiological

levels (Dahl et al., 2006; Hembree et al., 2009). For patients using intramuscular (IM) testosterone cypionate or enanthate, some clinicians check trough levels while others prefer midcycle levels (Dahl et al., 2006; Hembree et al., 2009; Tangpricha, Turner, Malabanan, & Holick, 2001; Tangpricha, Ducharme, Barber, & Chipkin, 2003).

Monitoring for adverse events should include both clinical and laboratory evaluation. Follow-up should include careful assessment for signs and symptoms of excessive weight gain, acne, uterine break-through bleeding, and cardiovascular impairment, as well as psychiatric symptoms in at-risk patients. Physical examinations should include measurement of pressure, weight, pulse, and skin; and heart and lung exams (Feldman & Safer, 2009). Laboratory monitoring should be based on the risks of hormone therapy described above, a patient's individual co-morbidities and risk factors, and the specific hormone regimen itself. Specific lab monitoring protocols have been published (Feldman & Safer, 2009; Hembree et al., 2009).

Hormone Regimens

To date, no controlled clinical trials of any feminizing/masculinizing hormone regimen have been conducted to evaluate safety or efficacy in producing physical transition. As a result, wide variation in doses and types of hormones have been published in the medical literature (Moore et al., 2003; Tangpricha et al., 2003; van Kesteren, Asscheman, Megens, & Gooren, 1997). In addition, access to particular medications may be limited by a patient's geographical location and/or social or economic situations. For these reasons, WPATH does not describe or endorse a particular feminizing/masculinizing hormone regimen. Rather, the medication classes and routes of administration used in most published regimens are broadly reviewed.

As outlined above, there are demonstrated safety differences in individual elements of various regimens. The Endocrine Society Guidelines (Hembree et al., 2009) and Feldman and Safer (2009) provide specific guidance regarding the types of hormones and suggested dosing to maintain levels within physiologic ranges for a patient's desired gender expression (based on goals of full feminization/masculinization). It is strongly recommend that hormone providers regularly review the literature for new information and use those medications that safely meet individual patient needs with available local resources.

Regimens for feminizing hormone therapy (MtF)

Estrogen

Use of oral estrogen, and specifically ethinyl estradiol, appears to increase the risk of VTE. Because of this safety concern, ethinyl estradiol is not recommended for feminizing hormone therapy. Transdermal estrogen is recommended for those patients with risks factors for VTE. The risk of adverse events increases with higher doses, particular those resulting in supraphysiologic levels (Hembree et al., 2009). Patients with co-morbid conditions that can be affected by estrogen should avoid oral estrogen if possible and be started at lower levels. Some patients may not be able to safely use the levels of estrogen needed to get the desired results. This possibility needs to be discussed with patients well in advance of starting hormone therapy.

Androgen reducing medications (“anti-androgens”)

A combination of estrogen and “anti-androgens” is the most commonly studied regimen for feminization. Androgen reducing medications, from a variety of classes of drugs, have the effect of reducing either endogenous testosterone levels or testosterone activity, and thus diminishing masculine characteristics such as body hair. They minimize the dosage of estrogen needed to suppress testosterone, thereby reducing the risks associated with high-dose exogenous estrogen (Prior, Vigna, Watson, Diewold, & Robinow, 1986; Prior, Vigna, & Watson, 1989).

Common anti-androgens include the following:

- Spironolactone, an antihypertensive agent, directly inhibits testosterone secretion and androgen binding to the androgen receptor. Blood pressure and electrolytes need to be monitored because of the potential for hyperkalemia.
- Cyproterone acetate is a progestational compound with anti-androgenic properties. This medication is not approved in the United States because of concerns over potential hepatotoxicity, but it is widely used elsewhere (De Cuypere et al., 2005).
- GnRH agonists (e.g., goserelin, buserelin, triptorelin) are neurohormones that block the gonadotropin releasing hormone receptor, thus blocking the release of follicle stimulating hormone and luteinizing hormone. This leads to highly effective gonadal blockade. However, these medications are expensive and only available as injectables or implants.
- 5-alpha reductase inhibitors (finasteride and dutasteride) block the conversion of testosterone to the more active agent, 5-alpha-dihydrotestosterone. These medications have beneficial effects on scalp hair loss, body hair growth, sebaceous glands, and skin consistency.

Cyproterone and spironolactone are the most commonly used anti-androgens and are likely the most cost-effective.

Progestins

With the exception of cyproterone, the inclusion of progestins in feminizing hormone therapy is controversial (Oriol, 2000). Because progestins play a role in mammary development on a cellular level, some clinicians believe that these agents are necessary for full breast development (Basson & Prior, 1998; Oriol, 2000). However, a clinical comparison of feminization regimens with and without progestins found that the addition of progestins neither enhanced breast growth nor lowered serum levels of free testosterone (Meyer III et al., 1986). There are concerns regarding potential adverse effects of progestins, including depression, weight gain, and lipid changes (Meyer III et al., 1986; Tangpricha et al., 2003). Progestins (especially medroxyprogesterone) are also suspected to increase breast cancer risk and cardiovascular risk in women (Rossouw et al., 2002). Micronized progesterone may be better tolerated and have a more favorable impact on the lipid profile than medroxyprogesterone does (de Lignières, 1999; Fitzpatrick, Pace, & Wiita, 2000).

Regimens for masculinizing hormone therapy (FtM)

Testosterone

Testosterone generally can be given orally, transdermally, or parenterally (IM), although buccal and implantable preparations are also available. Oral testosterone undecanoate, available outside the United States, results in lower serum testosterone levels than non-oral preparations and has limited efficacy in suppressing menses (Feldman, 2005, April; Moore et al., 2003). Because intramuscular testosterone cypionate or enanthate are often administered every 2-4 weeks, some patients may notice cyclic variation in effects (e.g., fatigue and irritability at the end of the injection cycle, aggression or expansive mood at the beginning of the injection cycle), as well as more time outside the normal physiologic levels (Jockenhövel, 2004). This may be mitigated by using a lower but more frequent dosage schedule or by using a daily transdermal preparation (Dobs et al., 1999; Jockenhövel, 2004; Nieschlag et al., 2004). Intramuscular testosterone undecanoate (not currently available in the United States) maintains stable, physiologic testosterone levels over approximately 12 weeks and has been effective in both the setting of hypogonadism and in FtM individuals (Mueller, Kiesewetter, Binder, Beckmann, & Dittrich, 2007; Zitzmann, Saad, & Nieschlag, 2006). There is evidence that transdermal and intramuscular testosterone achieve similar masculinizing results, although the timeframe may be somewhat slower with transdermal preparations (Feldman, 2005, April). Especially as patients age, the goal is to use the lowest dose needed to maintain the desired clinical result, with appropriate precautions being made to maintain bone density.

Other agents

Progestins, most commonly medroxyprogesterone, can be used for a short period of time to assist with menstrual cessation early in hormone therapy. GnRH agonists can be used similarly, as well as for refractory uterine bleeding in patients without an underlying gynecological abnormality.

Bioidentical and compounded hormones

As discussion surrounding the use of bioidentical hormones in postmenopausal hormone replacement has heightened, interest has also increased in the use of similar compounds in feminizing/masculinizing hormone therapy. There is no evidence that custom compounded bioidentical hormones are safer or more effective than government agency-approved bioidentical hormones (Sood, Shuster, Smith, Vincent, & Jatoi, 2011). Therefore, it has been advised by the North American Menopause Society (2010) and others to assume that, whether the hormone is from a compounding pharmacy or not, if the active ingredients are similar, it should have a similar side-effect profile. WPATH concurs with this assessment.

IX

Reproductive Health

Many transgender, transsexual, and gender nonconforming people will want to have children. Because feminizing/masculinizing hormone therapy limits fertility (Darney, 2008; Zhang, Gu, Wang, Cui, & Bremner, 1999), it is desirable for patients to make decisions concerning fertility before starting hormone therapy or undergoing surgery to remove/alter their reproductive organs. Cases are known of people who received hormone therapy and genital surgery and later regretted their inability to parent genetically related children (De Sutter, Kira, Verschoor, & Hotimsky, 2002).

Health care professionals – including mental health professionals recommending hormone therapy or surgery, hormone-prescribing physicians, and surgeons – should discuss reproductive options with patients prior to initiation of these medical treatments for gender dysphoria. These discussions should occur even if patients are not interested in these issues at the time of treatment, which may be more common for younger patients (De Sutter, 2009). Early discussions are desirable, but not always possible. If an individual has not had complete sex reassignment surgery, it may be possible to stop hormones long enough for natal hormones to recover, allowing the production of mature